
For each rational function in 1-5,

- Find the power function it most closely resembles for very large values of x.
- Describe the **long run** behavior by completing the boxes:

As
$$x \to -\infty$$
, then $y \to -\infty$; as $x \to \infty$, then $y \to -\infty$

c. Sketch the power function which has the same **long run** behavior. Pick from these choices: The *short run* behavior is covered up to emphasize that only the **long run** behavior is being mirrored.



d. Find the horizontal asymptote, if there is one. If none, state so.

1.
$$f(x) = \underbrace{\frac{8x^3}{+5x-9}}_{4x^7+200x^2-6}$$
 a. Power function model: $y = \underbrace{\frac{8x^3}{4x^7}}_{2x^4} = \underbrace{\frac{2}{x^4}}_{2x^4}$ (simplify) b. As $x \to -\infty$, then $y \to \boxed{0}$; as $x \to \infty$, then $y \to \boxed{0}$

a. Power function model:
$$y = \frac{\delta x}{4x^7} = \frac{2}{x^4}$$
 (simplify)

b. As
$$x \to -\infty$$
, then $y \to \boxed{0}$; as $x \to \infty$, then $y \to \boxed{0}$

- c. Long run behavior looks like this power function
- d. horizontal asymptote: y = 0



2.
$$f(x) = \frac{36x^3 + 3x - 7}{x^2 - 4x^3}$$

a. Power function model:
$$y = \frac{36x^3}{-4x^3} = -9$$
 (simplify)

b. As
$$x \to -\infty$$
, then $y \to \boxed{-9}$; as $x \to -\infty$, then $y \to \boxed{-9}$

d. horizontal asymptote:
$$y = -9$$



3.
$$f(x) = \frac{3 + 4x}{2 + 7x}$$

a. Power function model:
$$y = \frac{4x}{7x} = \frac{4}{7}$$
 (simplify)

b. As
$$x \to -\infty$$
, then $y \to \boxed{\frac{4}{7}}$; as $x \to \infty$, then $y \to \boxed{\frac{4}{7}}$

d. horizontal asymptote:
$$y = \frac{4}{7}$$



4.
$$f(x) = \frac{10x^6 - 4x}{(x-3)(x-4)}$$

= $\frac{10x^6 - 4x}{(x^2 + \text{remaining terms})}$

a. Power function model:
$$y = \frac{10x}{x^2} = 10x^4$$
 (simplify)

b. As
$$x \to -\infty$$
, then $y \to \boxed{\infty}$; as $x \to \infty$, then $y \to \boxed{\infty}$

5.
$$f(x) = \frac{2(x-2)^2(x-6)}{9(x-5)^3}$$

a. Power function model:
$$y = \frac{2x^3}{9x^3} = \frac{2}{9}$$
 (simplify)
b. As $x \to -\infty$, then $y \to \begin{bmatrix} \frac{2}{9} \end{bmatrix}$; as $x \to \infty$, then $y \to \infty$

$$= \frac{2x^3 + \text{remaining terms}}{9x^3 + \text{remaining terms}}$$

d. horizontal asymptote:
$$y = \frac{2}{9}$$



For the functions below, report the horizontal asymptote, if there is one. If none, state so.

6.
$$f(x) = \frac{7(x+2)(x+5)}{11(x-5)}$$
 9. $f(x) = \frac{6x}{3x^2+10} + 2$

9.
$$f(x) = \frac{6x}{3x^2 + 10} + 2$$

7.
$$f(x) = \frac{12x^2 + 1}{3x^2 + 2} + 3$$

10.
$$f(x) = \frac{3(x+5)^2(x-4)}{4(x-6)^3(x-1)}$$

7.
$$y = 7$$

8. $y = 1250$

8.
$$f(x) = \frac{25x^2 + 38}{x(1+0.02x)}$$

10.
$$f(x) = \frac{3(x+5)^2(x-4)}{4(x-6)^3(x-1)}$$

9.
$$y = 12$$

10.
$$y = 0$$

Please see next page for more

DETAILED SOLUTIONS

Easter Egg! For 1 bonus participation point email John LaMaster, lamaster@pfw.edu, by 11:59 PM April 16 to share the name of the monster that makes a brief cameo appearance.

6. $f(x) = \frac{7(x+2)(x+5)}{11(x-5)}$ looks like the power function $\frac{7x^2}{11x} = \frac{7x}{11}$ for very large values of x.

This is a line of positive slope. There is no horizontal line this function approaches.

As $x \to -\infty$, then $y \to -\infty$; as $x \to \infty$, then $y \to \infty$.

So, no horizontal asymptote.

7. $f(x) = \frac{12x^2 + 1}{3x^2 + 2} + 3$ is a vertical shift of the function $y = \frac{12x^2 + 1}{3x^2 + 2}$ up 3.

Since $y = \frac{12x^2 + 1}{3x^2 + 2}$ looks like the power function $\frac{12x^2}{3x^2} = 4$ for very large values of x,

the graph of $y = \frac{12x^2 + 1}{3x^2 + 2}$ would have a horizontal asymptote of y = 4. Therefore, the shifted

function $f(x) = \frac{12x^2 + 1}{3x^2 + 2} + 3$ would have a horizontal asymptote of y = 7. Note that for very

large values of x, $f(x) = \frac{12x^2 + 1}{3x^2 + 2} + 3 \implies \frac{12x^2}{3x^2} + 3 = 4 + 3 = 7$.

- 8. $f(x) = \frac{25x^2 + 38}{x(1 + 0.02x)}$ looks like the power function $\frac{25x^2}{0.02x^2} = \frac{25}{0.02} = 1250$ for very large values of x. Therefore, it has a horizontal asymptote of y = 1250.
- 9. $f(x) = \frac{6x}{3x^2 + 10} + 2$ is a vertical shift of the function $y = \frac{6x}{3x^2 + 10}$ up 2.

Since $y = \frac{6x}{3x^2 + 10}$ looks like the power function $\frac{6x}{3x^2} = \frac{2}{x}$ for very large values of x,

the graph of $y = \frac{6x}{3x^2 + 10}$ would have a horizontal asymptote of y = 0. Therefore, the shifted

function $f(x) = \frac{6x}{3x^2 + 10} + 2$ would have a horizontal asymptote of y = 2. Note that for very

large values of x, $\frac{6x}{3x^2} + 2 = \frac{2}{x} + 2 \rightarrow 0 + 2$

10. $f(x) = \frac{3(x+5)^2(x-4)}{4(x-6)^3(x-1)} = \frac{3x^3 + \text{remaining terms}}{4x^4 + \text{remaining terms}}$, so it looks like the power function $\frac{3x^3}{4x^4}$

for very large values of x. Note that $\frac{3x^3}{4x^4} = \frac{3}{4x}$ approaches 0 as x increases without bound, so

 $f(x) = \frac{3(x+5)^2(x-4)}{4(x-6)^3(x-1)}$ has a horizontal asymptote of y = 0.