

UNIQUE CONTINUATION FOR A GRADIENT INEQUALITY WITH L^n POTENTIAL

ADAM COFFMAN, YIFEI PAN, AND YUAN ZHANG

ABSTRACT. We establish a unique continuation property for solutions of the differential inequality $|\nabla u| \leq V|u|$, where V is locally L^n integrable on a domain in \mathbb{R}^n . A stronger uniqueness result is obtained if in addition the solutions are locally Lipschitz. One application is a finite order vanishing property in the L^2 sense for the exponential of $W^{1,n}$ functions. We further discuss related results for the Cauchy-Riemann operator $\bar{\partial}$ and characterize the vanishing order for smooth extension of holomorphic functions across the boundary.

1. INTRODUCTION AND RESULTS

Let Ω be a connected open subset of \mathbb{R}^n . We investigate solutions to the following differential inequality concerning the gradient operator ∇ :

$$|\nabla u| \leq V|u| \quad \text{on } \Omega, \quad (1.1)$$

with the potential $V \in L^n_{loc}(\Omega)$.

A function $u \in L^2_{loc}(\Omega)$ is said to vanish to infinite order (or to be flat) at a point $x_0 \in \Omega$ (in the L^2 sense) means that for all $m \geq 0$,

$$\lim_{r \rightarrow 0} r^{-m} \int_{|x-x_0| < r} |u(x)|^2 dv = 0,$$

where dv is the Lebesgue measure element in \mathbb{R}^n . Otherwise, u vanishes to finite order at x_0 in the L^2 sense. We say a differential (in)equality satisfies the (strong) unique continuation property to mean that every $H^1_{loc}(\Omega)$ ($= W^{1,2}_{loc}(\Omega)$) solution that vanishes to infinite order at a point in the L^2 sense must vanish identically. Here for $p \geq 1$, $W^{1,p}_{loc}(\Omega)$ is the standard Sobolev space of $L^p_{loc}(\Omega)$ functions whose first order weak derivatives are represented by functions in $L^p_{loc}(\Omega)$. While studying the unique continuation property of the Cauchy-Riemann operator $\bar{\partial}$ in several complex variables:

$$|\bar{\partial} u| \leq V|u| \quad \text{on } \Omega \subseteq \mathbb{C}^n, \quad (1.2)$$

we observe that (1.2) is reduced to (1.1) when the solutions are real-valued. This motivates us to study the following unique continuation property of $H^1_{loc}(\Omega)$ solutions to (1.1).

2020 *Mathematics Subject Classification.* Primary 35R45; Secondary 26B35, 32A40, 35A02.

Key words and phrases. unique continuation, differential inequalities, divergent integrals.

Theorem 1.1. *Let Ω be a domain in \mathbb{R}^n , $n \geq 2$ and $V \in L^n_{loc}(\Omega)$. Suppose $u = (u_1, \dots, u_M) : \Omega \rightarrow \mathbb{R}^M$ with $u \in H^1_{loc}(\Omega)$ and satisfies $|\nabla u| \leq V|u|$ a.e. on Ω . If u vanishes to infinite order at some $x_0 \in \Omega$, then $u \equiv 0$.*

The $n = 2$ case in Theorem 1.1 is due to a unique continuation property result in [PZ] concerning the $\bar{\partial}$ operator. For higher dimensions, the proof makes use of a Hardy-type inequality, along with the Gagliardo-Nirenberg-Sobolev inequality. When the potential is no longer in L^n , one can still get the unique continuation property for some special types of potentials, see Theorem 2.6. However, as shown in Example 2.5, the property fails in general for $V \notin L^n$. On the other hand, Theorem 2.7 states that the weak unique continuation property always holds for (1.1) as long as $V \in L^2$.

As a consequence of Theorem 1.1, we obtain the following property of vanishing to finite order for the exponential of $W^{1,n}$ functions. Note that the $W^{1,n}$ space is the critical Sobolev space where the Sobolev embedding theorem fails, and instead is substituted by the Moser-Trudinger inequality.

Theorem 1.2. *Let Ω be an open set in \mathbb{R}^n , $n \geq 2$. Suppose $\phi : \Omega \rightarrow \mathbb{R}$ with $\phi \in W^{1,n}_{loc}(\Omega)$. Then the exponential e^ϕ of ϕ vanishes to finite order in the L^2 sense at each point in Ω .*

In the second part of the paper, we focus on locally Lipschitz solutions to (1.1). Under the context of this more restricted function space, we are able to prove a uniqueness result below by just assuming the vanishing of the first jets. A similar uniqueness result was discussed in [PW] for higher order differential operators on smooth functions of one variable ($n = 1$). It is worth pointing out that the Lipschitz assumption on the solutions cannot be dropped here when $n \geq 2$, see Remark 3.9.

Theorem 1.3. *Let Ω be a domain in \mathbb{R}^n , $n \geq 1$ and $V \in L^n_{loc}(\Omega)$. Suppose $u = (u_1, \dots, u_M) : \Omega \rightarrow \mathbb{R}^M$ is a locally Lipschitz function on Ω satisfying $|\nabla u| \leq V|u|$ a.e. on Ω . If $u(x_0) = 0$ at some $x_0 \in \Omega$, then $u \equiv 0$.*

Theorem 1.3 can be readily applied to study the uniqueness of some types of nonlinear differential systems, as indicated in Corollary 3.11. In Section 4, we discuss further applications under the Lipschitz setting. In particular, Theorem 4.2 shows that the logarithm of a positive Lipschitz function cannot fall in $W^{1,n}$ near every zero point of the function. On the other hand, we prove that if in addition e^ϕ in Theorem 1.2 is Lipschitz, then e^ϕ must be nowhere zero, see Corollary 4.5.

In the last section, we discuss related results for the $\bar{\partial}$ operator on domains in \mathbb{C}^n . To start with, we construct Example 5.1 to show that the gradient operator ∇ in Theorem 1.3 cannot be replaced by the $\bar{\partial}$ operator even for real analytic functions. On the other hand, we give finer characterizations in terms of an L^2 divergence for holomorphic functions that are extended smoothly across the boundary.

2. UNIQUE CONTINUATION FOR H^1 SOLUTIONS

Let Ω be a domain (by which we mean a connected open set) in $\mathbb{R}^n = \{(x_1, \dots, x_n)\}$. For scalar valued $u : \Omega \rightarrow \mathbb{R}$, $u \in W_{loc}^{1,p}(\Omega)$, the gradient of u is the vector of first order weak partial derivatives:

$$\nabla u = (\partial_{x_1} u, \dots, \partial_{x_n} u),$$

defined on Ω . The norm of a vector $x \in \mathbb{R}^n$ is $|x| = \sqrt{x_1^2 + \dots + x_n^2}$, and in particular the norm of the gradient is defined on the domain of the gradient by

$$|\nabla u|^2 = (\partial_{x_1} u)^2 + \dots + (\partial_{x_n} u)^2.$$

Given $r > 0$ and $x_0 \in \mathbb{R}^n$, let $B_r(x_0)$ denote the open ball centered at x_0 with radius r — in the special case $x_0 = 0$, we abbreviate $B_r = B_r(0)$. For sets $A \subseteq \mathbb{R}^n$, let ∂A denote the set of boundary points of A and let \bar{A} denote the closure of A .

In this section, we prove Theorem 1.1, the unique continuation property for vector valued H^1 solutions $u : \Omega \rightarrow \mathbb{R}^M$, where the inequality (1.1) reads as

$$|\nabla u| = \left(\sum_{j=1}^n \sum_{k=1}^M |\partial_{x_j} u_k|^2 \right)^{\frac{1}{2}} \leq V \left(\sum_{k=1}^M |u_k|^2 \right)^{\frac{1}{2}} = V|u|. \quad (2.1)$$

We first prove the following Hardy-type inequality for ∇ .

Lemma 2.1. *Let $u \in H^1(\mathbb{R}^n)$ with support outside a neighborhood of 0. Then for any $\lambda > \frac{n}{2}$,*

$$\int_{\mathbb{R}^n} \frac{|u(x)|^2}{|x|^{2\lambda}} dv \leq \frac{4}{(2\lambda - n)^2} \int_{\mathbb{R}^n} \frac{|\nabla u(x)|^2}{|x|^{2\lambda-2}} dv. \quad (2.2)$$

Proof. We first show the inequality when $u \in C_c^\infty(\mathbb{R}^n \setminus \{0\})$. Let

$$F(x) := \sum_{j=1}^n \frac{|u(x)|^2 x_j}{|x|^{2\lambda}} dx_1 \wedge \dots \wedge dx_{j-1} \wedge dx_{j+1} \wedge \dots \wedge dx_n.$$

Then F is a smooth $(n-1)$ form with compact support. Note for each $j = 1, \dots, n$,

$$\sum_{j=1}^n \partial_{x_j} \left(\frac{x_j}{|x|^{2\lambda}} \right) = \sum_{j=1}^n \left(\frac{1}{|x|^{2\lambda}} - \frac{2\lambda x_j^2}{|x|^{2\lambda+2}} \right) = \frac{n - 2\lambda}{|x|^{2\lambda}}.$$

Applying Stokes' theorem on F , we have

$$0 = \int_{\mathbb{R}^n} dF = \int_{\mathbb{R}^n} \frac{(n - 2\lambda)|u(x)|^2}{|x|^{2\lambda}} dv + \int_{\mathbb{R}^n} \frac{2u(x)\langle \nabla u(x), x \rangle}{|x|^{2\lambda}} dv.$$

Thus

$$\int_{\mathbb{R}^n} \frac{|u(x)|^2}{|x|^{2\lambda}} dv = \frac{2}{(2\lambda - n)} \int_{\mathbb{R}^n} \frac{u(x)\langle \nabla u(x), x \rangle}{|x|^{2\lambda}} dv.$$

By the Cauchy-Schwarz inequality, one further gets

$$\begin{aligned} \int_{\mathbb{R}^n} \frac{|u(x)|^2}{|x|^{2\lambda}} dv &\leq \frac{2}{(2\lambda - n)} \int_{\mathbb{R}^n} \frac{|u(x)| |\nabla u(x)|}{|x|^{2\lambda-1}} dv \\ &\leq \frac{2}{(2\lambda - n)} \left(\int_{\mathbb{R}^n} \frac{|u(x)|^2}{|x|^{2\lambda}} dv \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}^n} \frac{|\nabla u(x)|^2}{|x|^{2\lambda-2}} dv \right)^{\frac{1}{2}}. \end{aligned}$$

Dividing both sides by $\left(\int_{\mathbb{R}^n} \frac{|u(x)|^2}{|x|^{2\lambda}} dv \right)^{\frac{1}{2}}$ and then squaring both sides, we obtain (2.2) for $u \in C_c^\infty(\mathbb{R}^n \setminus \{0\})$.

For general $u \in H^1(\mathbb{R}^n)$ with support, say, away from $B_r, r > 0$, we use the standard density argument. In detail, let $u^{(j)} \in C_c^\infty(\mathbb{R}^n \setminus B_r) \rightarrow u$ in H^1 norm. Then

$$\begin{aligned} \left(\int_{\mathbb{R}^n} \frac{|u(x)|^2}{|x|^{2\lambda}} dv \right)^{\frac{1}{2}} &\leq \left(\int_{\mathbb{R}^n \setminus B_r} \frac{|u(x) - u^{(j)}(x)|^2}{|x|^{2\lambda}} dv \right)^{\frac{1}{2}} + \left(\int_{\mathbb{R}^n} \frac{|u^{(j)}(x)|^2}{|x|^{2\lambda}} dv \right)^{\frac{1}{2}} \\ &\leq \frac{1}{r^\lambda} \left(\int_{\mathbb{R}^n} |u(x) - u^{(j)}(x)|^2 dv \right)^{\frac{1}{2}} + \frac{2}{(2\lambda - n)} \left(\int_{\mathbb{R}^n} \frac{|\nabla u^{(j)}(x)|^2}{|x|^{2\lambda-2}} dv \right)^{\frac{1}{2}}. \end{aligned}$$

Here we used (2.2) for $u^{(j)} \in C_c^\infty(\mathbb{R}^n \setminus \{0\})$. Thus

$$\begin{aligned} &\left(\int_{\mathbb{R}^n} \frac{|u(x)|^2}{|x|^{2\lambda}} dv \right)^{\frac{1}{2}} \\ &\leq \frac{1}{r^\lambda} \left(\int_{\mathbb{R}^n} |u(x) - u^{(j)}(x)|^2 dv \right)^{\frac{1}{2}} + \frac{2}{(2\lambda - n)} \left(\int_{\mathbb{R}^n \setminus B_r} \frac{|\nabla u^{(j)}(x) - \nabla u(x)|^2}{|x|^{2\lambda-2}} dv \right)^{\frac{1}{2}} \\ &\quad + \frac{2}{(2\lambda - n)} \left(\int_{\mathbb{R}^n} \frac{|\nabla u(x)|^2}{|x|^{2\lambda-2}} dv \right)^{\frac{1}{2}} \\ &\leq \frac{1}{r^\lambda} \left(\int_{\mathbb{R}^n} |u(x) - u^{(j)}(x)|^2 dv \right)^{\frac{1}{2}} + \frac{2}{(2\lambda - n)r^{\lambda-1}} \left(\int_{\mathbb{R}^n} |\nabla u^{(j)}(x) - \nabla u(x)|^2 dv \right)^{\frac{1}{2}} \\ &\quad + \frac{2}{(2\lambda - n)} \left(\int_{\mathbb{R}^n} \frac{|\nabla u(x)|^2}{|x|^{2\lambda-2}} dv \right)^{\frac{1}{2}} \\ &\leq \left(\frac{1}{r^\lambda} + \frac{2}{(2\lambda - n)r^{\lambda-1}} \right) \|u - u^{(j)}\|_{H^1(\mathbb{R}^n)} + \frac{2}{(2\lambda - n)} \left(\int_{\mathbb{R}^n} \frac{|\nabla u(x)|^2}{|x|^{2\lambda-2}} dv \right)^{\frac{1}{2}}. \end{aligned}$$

Letting $j \rightarrow \infty$, we have the desired inequality (2.2). ■

Lemma 2.2. *Let $u \in H^1(\mathbb{R}^n)$ with support outside a neighborhood of 0. Then there exists a constant $C_0 > 0$ such that for any $\lambda \gg \frac{n}{2}$,*

$$\int_{\mathbb{R}^n} \left| \nabla \left(\frac{u(x)}{|x|^{\lambda-1}} \right) \right|^2 dv \leq C_0 \int_{\mathbb{R}^n} \frac{|\nabla u(x)|^2}{|x|^{2\lambda-2}} dv.$$

Proof. Since $|\nabla|x|| = 1$, we have

$$\begin{aligned} \int_{\mathbb{R}^n} \left| \nabla \left(\frac{u(x)}{|x|^{\lambda-1}} \right) \right|^2 dv &\leq 2 \int_{\mathbb{R}^n} \frac{|\nabla u(x)|^2}{|x|^{2\lambda-2}} dv + 2(\lambda-1)^2 \int_{\mathbb{R}^n} \frac{|u(x)|^2}{|x|^{2\lambda}} dv \\ &\leq 2 \int_{\mathbb{R}^n} \frac{|\nabla u(x)|^2}{|x|^{2\lambda-2}} dv + \frac{8(\lambda-1)^2}{(2\lambda-n)^2} \int_{\mathbb{R}^n} \frac{|\nabla u(x)|^2}{|x|^{2\lambda-2}} dv \\ &= \left(2 + \frac{8(\lambda-1)^2}{(2\lambda-n)^2} \right) \int_{\mathbb{R}^n} \frac{|\nabla u(x)|^2}{|x|^{2\lambda-2}} dv. \end{aligned}$$

Here in the second inequality we used Lemma 2.1. The lemma thus follows from the fact that $\lim_{\lambda \rightarrow \infty} \frac{8(\lambda-1)^2}{(2\lambda-n)^2} = 2$. \blacksquare

Throughout the rest of the paper, we occasionally use the notation $a \lesssim b$ for two quantities a and b , to mean that there exists a universal constant C (dependent only possibly on n) such that $a \leq Cb$. To prove Theorem 1.1 in the case when $n = 2$, we will use the following unique continuation property established in [PZ] for $\bar{\partial}$. Note that identifying $z \in \mathbb{C}$ with $(x_1, x_2) \in \mathbb{R}^2$, then for a function u on Ω , $\bar{\partial}_z u = \frac{1}{2}(\partial_{x_1} u + i\partial_{x_2} u)$. It would be interesting to have a real-variable approach for this case, but we currently do not.

Proposition 2.3. [PZ] *Let Ω be a domain in \mathbb{C} . Suppose $u = (u_1, \dots, u_N) : \Omega \rightarrow \mathbb{C}^N$ with $u \in H_{loc}^1(\Omega)$ and satisfies $|\bar{\partial}u| \leq V|u|$ a.e. on Ω for some $V \in L_{loc}^2(\Omega)$. If u vanishes to infinite order at $z_0 \in \Omega$, then u vanishes identically.* \blacksquare

Proof of Theorem 1.1: The $n = 2$ case follows from Proposition 2.3 and the trivial fact that $|\bar{\partial}u| \lesssim |\nabla u|$. When $n \geq 3$, without loss of generality assume $x_0 = 0$. Fix $r \in (0, 1)$ so that

$$\left(\int_{B_{2r}} |V(x)|^n dv \right)^{\frac{2}{n}} < \frac{1}{2C_0 C_1^2}, \quad (2.3)$$

where C_0 is the constant in Lemma 2.2, and C_1 is the constant in the Gagliardo-Nirenberg-Sobolev inequality:

$$\|f\|_{L^{\frac{2n}{n-2}}(\mathbb{R}^n)} \leq C_1 \|\nabla f\|_{L^2(\mathbb{R}^n)}, \quad \text{for all } f \in H^1(\mathbb{R}^n).$$

We shall show that $u = 0$ in $B_{\frac{r}{2}}$. Thus, applying a standard propagation argument we obtain $u \equiv 0$ on Ω .

Choose $\eta \in C_c^\infty(\mathbb{R}^n)$ such that $0 \leq \eta \leq 1$, $\eta = 1$ on B_r , $\eta = 0$ outside B_{2r} , and $|\nabla \eta| \leq \frac{2}{r}$ on $B_{2r} \setminus B_r$. Let $\psi \in C^\infty(\mathbb{R}^n)$ be such that $0 \leq \psi \leq 1$, $\psi = 0$ in B_1 , $\psi = 1$ outside B_2 , and

$|\nabla\psi| \leq 2$ on $B_2 \setminus B_1$. For each $k \geq \frac{4}{r}$ (then $\frac{2}{k} \leq \frac{r}{2}$), let $\psi_k(x) = \psi(kx)$, $x \in \mathbb{R}^n$. Defining $u^{(k)} = \psi_k \eta u$, note that $u^{(k)} \in H^1(\mathbb{R}^n)$ and is supported inside $B_{2r} \setminus B_{\frac{1}{k}}$. Then for each $k \geq \frac{4}{r}$ and $\lambda > \frac{n}{2}$,

$$\begin{aligned}
& \int_{B_{2r}} \frac{|\nabla u^{(k)}(x)|^2}{|x|^{2\lambda-2}} dv \\
& \lesssim \int_{B_{2r}} \frac{|\psi_k(x)\eta(x)|^2 |\nabla u(x)|^2}{|x|^{2\lambda-2}} dv + \int_{B_r} \frac{|\nabla \psi_k(x)|^2 |u(x)|^2}{|x|^{2\lambda-2}} dv + \int_{B_{2r} \setminus B_r} \frac{|\nabla \eta(x)|^2 |u(x)|^2}{|x|^{2\lambda-2}} dv \\
& \leq \int_{B_{2r}} \frac{|V(x)|^2 |\psi_k(x)\eta(x)u(x)|^2}{|x|^{2\lambda-2}} dv + \int_{B_r} \frac{|\nabla \psi_k(x)|^2 |u(x)|^2}{|x|^{2\lambda-2}} dv + \int_{B_{2r} \setminus B_r} \frac{|\nabla \eta(x)|^2 |u(x)|^2}{|x|^{2\lambda-2}} dv \\
& \leq \left(\int_{B_{2r}} |V(x)|^n dv \right)^{\frac{2}{n}} \left(\int_{\mathbb{R}^n} \left(\frac{|u^{(k)}(x)|}{|x|^{\lambda-1}} \right)^{\frac{2n}{n-2}} dv \right)^{\frac{n-2}{n}} + \int_{B_r} \frac{|\nabla \psi_k(x)|^2 |u(x)|^2}{|x|^{2\lambda-2}} dv \\
& \quad + \int_{B_{2r} \setminus B_r} \frac{|\nabla \eta(x)|^2 |u(x)|^2}{|x|^{2\lambda-2}} dv. \tag{2.4}
\end{aligned}$$

Here we have used Hölder's inequality in (2.4). Since $\frac{|u^{(k)}(x)|}{|x|^{\lambda-1}} \in H^1(\mathbb{R}^n)$, $n \geq 3$, making use of the Gagliardo-Nirenberg-Sobolev inequality and Lemma 2.2, we get

$$\left(\int_{\mathbb{R}^n} \left(\frac{|u^{(k)}(x)|}{|x|^{\lambda-1}} \right)^{\frac{2n}{n-2}} dv \right)^{\frac{n-2}{n}} \leq C_1^2 \int_{\mathbb{R}^n} \left| \nabla \left(\frac{|u^{(k)}(x)|}{|x|^{\lambda-1}} \right) \right|^2 dv \leq C_0 C_1^2 \int_{B_{2r}} \frac{|\nabla u^{(k)}(x)|^2}{|x|^{2\lambda-2}} dv.$$

This combined with (2.4) and (2.3) for each $k \geq \frac{4}{r}$ and $\lambda > \frac{n}{2}$ leads to

$$\int_{B_{2r}} \frac{|\nabla u^{(k)}(x)|^2}{|x|^{2\lambda-2}} dv \leq 2 \int_{B_r} \frac{|\nabla \psi_k(x)|^2 |u(x)|^2}{|x|^{2\lambda-2}} dv + 2 \int_{B_{2r} \setminus B_r} \frac{|\nabla \eta(x)|^2 |u(x)|^2}{|x|^{2\lambda-2}} dv. \tag{2.5}$$

Now suppose toward a contradiction that $\nabla u \not\equiv 0$ on $B_{\frac{r}{2}}$. Then there exists $k_1 > 0$ such that

$$M_1 = \int_{B_{\frac{r}{2}} \setminus B_{\frac{2}{k_1}}} |\nabla u(x)|^2 dv > 0. \tag{2.6}$$

Consequently for each fixed $\lambda > \frac{n}{2}$,

$$M_\lambda = \int_{B_{\frac{r}{2}} \setminus B_{\frac{2}{k_1}}} \frac{|\nabla u(x)|^2}{|x|^{2\lambda-2}} dv > 0.$$

Noting that $\nabla u^{(k)} = \nabla u$ on $B_{\frac{r}{2}} \setminus B_{\frac{2}{k_1}}$ for any $k \geq k_1$ by construction of $u^{(k)}$, we further have for any $k \geq k_1$,

$$\int_{B_{\frac{r}{2}}} |\nabla u^{(k)}(x)|^2 dv \geq \int_{B_{\frac{r}{2}} \setminus B_{\frac{2}{k_1}}} |\nabla u(x)|^2 dv = M_1 > 0, \quad (2.7)$$

and

$$\int_{B_{\frac{r}{2}}} \frac{|\nabla u^{(k)}(x)|^2}{|x|^{2\lambda-2}} dv \geq \int_{B_{\frac{r}{2}} \setminus B_{\frac{2}{k_1}}} \frac{|\nabla u(x)|^2}{|x|^{2\lambda-2}} dv = M_\lambda > 0. \quad (2.8)$$

On the other hand, by flatness of u at 0,

$$\int_{B_r} \frac{|\nabla \psi_k(x)|^2 |u(x)|^2}{|x|^{2\lambda-2}} dv = \int_{B_{\frac{2}{k}} \setminus B_{\frac{1}{k}}} \frac{|\nabla \psi_k(x)|^2 |u(x)|^2}{|x|^{2\lambda-2}} dv \leq 4k^{2\lambda} \int_{B_{\frac{2}{k}}} |u(x)|^2 dv \rightarrow 0 \quad (2.9)$$

as $k \rightarrow \infty$. In particular, by (2.8) one can get some $k_\lambda > k_1$ such that

$$\int_{B_r} \frac{|\nabla \psi_{k_\lambda}(x)|^2 |u(x)|^2}{|x|^{2\lambda-2}} dv \leq \frac{M_\lambda}{4} \leq \frac{1}{4} \int_{B_r} \frac{|\nabla u^{(k_\lambda)}(x)|^2}{|x|^{2\lambda-2}} dv.$$

Thus (2.5) with $k = k_\lambda$ becomes

$$\int_{B_{2r}} \frac{|\nabla u^{(k_\lambda)}(x)|^2}{|x|^{2\lambda-2}} dv \leq 4 \int_{B_{2r} \setminus B_r} \frac{|\nabla \eta(x)|^2 |u(x)|^2}{|x|^{2\lambda-2}} dv. \quad (2.10)$$

Since

$$\int_{B_{2r}} \frac{|\nabla u^{(k_\lambda)}(x)|^2}{|x|^{2\lambda-2}} dv \geq \int_{B_{\frac{r}{2}}} \frac{|\nabla u^{(k_\lambda)}(x)|^2}{|x|^{2\lambda-2}} dv \geq \left(\frac{2}{r}\right)^{2\lambda-2} \int_{B_{\frac{r}{2}}} |\nabla u^{(k_\lambda)}(x)|^2 dv$$

and

$$\int_{B_{2r} \setminus B_r} \frac{|\nabla \eta(x)|^2 |u(x)|^2}{|x|^{2\lambda-2}} dv \leq \frac{1}{r^{2\lambda-2}} \int_{B_{2r} \setminus B_r} |\nabla \eta(x)|^2 |u(x)|^2 dv,$$

we obtain from (2.10) that

$$2^{2\lambda-4} \int_{B_{\frac{r}{2}}} |\nabla u^{(k_\lambda)}(x)|^2 dv \leq \int_{B_{2r} \setminus B_r} |\nabla \eta(x)|^2 |u(x)|^2 dv.$$

Letting $\lambda \rightarrow \infty$ and making use of the fact that $u \in H_{loc}^1(\Omega)$, we see that

$$\int_{B_{\frac{r}{2}}} |\nabla u(x)|^2 dv = 0.$$

But this contradicts (2.6)! We thus have $\nabla u \equiv 0$ on $B_{\frac{r}{2}}$. By flatness of u at 0, u must be zero on $B_{\frac{r}{2}}$. ■

Example 2.4. Given $0 < \varepsilon < \frac{n-1}{n}, n \geq 2$, consider the differential equation

$$|\nabla u| = V|u| \quad \text{on } B_{\frac{1}{2}},$$

where

$$V = \frac{\varepsilon(-\log|x|)^{\varepsilon-1}}{|x|} \quad \text{on } B_{\frac{1}{2}}.$$

It is straightforward to verify that $V \in L^n(B_{\frac{1}{2}})$. As a consequence of Theorem 1.1, every nonconstant H^1 solution must vanish to finite order in the L^2 sense at each point in $B_{\frac{1}{2}}$.

On the other hand, the function

$$u_0(x) = e^{-(\log|x|)^\varepsilon}$$

(extended to the origin by $u_0(0) = 0$) is continuous on $B_{\frac{1}{2}}$, and smooth on $B_{\frac{1}{2}} \setminus \{0\}$. Moreover, $u_0 \in H^1(B_{\frac{1}{2}})$ and is a solution of $|\nabla u| = Vu$ a.e. on $B_{\frac{1}{2}}$. Note that there is no contradiction with Theorem 1.1 since u_0 vanishes to finite order in the L^2 sense everywhere in $B_{\frac{1}{2}}$.

When $V \in L^p$, $p < n$, the unique continuation property fails in general as seen below.

Example 2.5. For each $1 \leq p < n$, and $0 < \epsilon < \frac{n-p}{p}$ (so that $(\epsilon + 1)p < n$),

$$u(x) = e^{-\frac{1}{|x|^\epsilon}}$$

(extended to the origin by $u(0) = 0$) is a smooth function on B_1 and vanishes to infinite order at 0. Moreover, the function u satisfies $|\nabla u| \leq V|u|$ on B_1 with

$$V = \frac{\epsilon}{|x|^{\epsilon+1}} \in L^p(B_1).$$

On the other hand, the following theorem states that for some special potentials in the form of multiples of $\frac{1}{|x|}$, the unique continuation property can still hold. Note that $\frac{1}{|x|} \notin L_{loc}^n$.

Theorem 2.6. *Let Ω be a domain in $\mathbb{R}^n, n \geq 1$. Suppose $u = (u_1, \dots, u_M) : \Omega \rightarrow \mathbb{R}^M$ with $u \in H_{loc}^1(\Omega)$ and satisfies $|\nabla u| \leq \frac{C}{|x|}|u|$ a.e. for some constant $C > 0$. If u vanishes to infinite order at some $x_0 \in \Omega$ in the L^2 sense, then u vanishes identically.*

Proof. Assume $x_0 = 0$ and consider $u^{(k)} = \psi_k \eta u$, where ψ_k and η are defined as in the proof of Theorem 1.1. Then by Lemma 2.1,

$$\begin{aligned}
 & \int_{B_{2r}} \frac{|u^{(k)}(x)|^2}{|x|^{2\lambda}} dv \\
 & \leq \frac{4}{(2\lambda - n)^2} \int_{B_{2r}} \frac{|\nabla u^{(k)}(x)|^2}{|x|^{2\lambda-2}} dv \\
 & \lesssim \frac{4}{(2\lambda - n)^2} \int_{B_{2r}} \frac{|\psi_k(x)\eta(x)|^2 |\nabla u(x)|^2}{|x|^{2\lambda-2}} dv + \frac{4}{(2\lambda - n)^2} \int_{B_r} \frac{|\nabla \psi_k(x)|^2 |u(x)|^2}{|x|^{2\lambda-2}} dv \\
 & \quad + \frac{4}{(2\lambda - n)^2} \int_{B_{2r} \setminus B_r} \frac{|\nabla \eta(x)|^2 |u(x)|^2}{|x|^{2\lambda-2}} dv \\
 & \leq \frac{4C^2}{(2\lambda - n)^2} \int_{B_{2r}} \frac{|u^{(k)}(x)|^2}{|x|^{2\lambda}} dv + \frac{4}{(2\lambda - n)^2} \int_{B_r} \frac{|\nabla \psi_k(x)|^2 |u(x)|^2}{|x|^{2\lambda-2}} dv \\
 & \quad + \frac{4}{(2\lambda - n)^2} \int_{B_{2r} \setminus B_r} \frac{|\nabla \eta(x)|^2 |u(x)|^2}{|x|^{2\lambda-2}} dv.
 \end{aligned}$$

Here we used the inequality $|\nabla u| \leq \frac{C}{|x|}|u|$ in the first term of the last inequality. When $\frac{4C^2}{(2\lambda-n)^2} \leq \frac{1}{2}$ (equivalently, when $\lambda > \frac{n}{2} + \sqrt{2}C$), one can move this first term to the left hand side and get

$$\int_{B_{2r}} \frac{|u^{(k)}(x)|^2}{|x|^{2\lambda}} dv \leq \frac{8}{(2\lambda - n)^2} \int_{B_r} \frac{|\nabla \psi_k(x)|^2 |u(x)|^2}{|x|^{2\lambda-2}} dv + \frac{8}{(2\lambda - n)^2} \int_{B_{2r} \setminus B_r} \frac{|\nabla \eta(x)|^2 |u(x)|^2}{|x|^{2\lambda-2}} dv.$$

Letting $k \rightarrow \infty$ and making use of the flatness of u with a similar argument as in (2.9), we obtain

$$\int_{B_{2r}} \frac{|u(x)|^2}{|x|^{2\lambda}} dv \leq \frac{16}{(2\lambda - n)^2} \int_{B_{2r} \setminus B_r} \frac{|\nabla \eta(x)|^2 |u(x)|^2}{|x|^{2\lambda-2}} dv.$$

Since

$$\int_{B_{2r}} \frac{|u(x)|^2}{|x|^{2\lambda}} dv \geq \int_{B_{\frac{r}{2}}} \frac{|u(x)|^2}{|x|^{2\lambda}} dv \geq \left(\frac{2}{r}\right)^{2\lambda} \int_{B_{\frac{r}{2}}} |u(x)|^2 dv$$

and

$$\int_{B_{2r} \setminus B_r} \frac{|\nabla \eta(x)|^2 |u(x)|^2}{|x|^{2\lambda-2}} dv \leq \frac{1}{r^{2\lambda-2}} \int_{B_{2r} \setminus B_r} |\nabla \eta(x)|^2 |u(x)|^2 dv,$$

we have

$$\int_{B_{\frac{r}{2}}} |u(x)|^2 dv \leq \frac{r^2}{(2\lambda - n)^2 2^{2\lambda-4}} \int_{B_{2r} \setminus B_r} |\nabla \eta(x)|^2 |u(x)|^2 dv.$$

Letting $\lambda \rightarrow \infty$, we see $u \equiv 0$ on $B_{\frac{r}{2}}$. ■

Although the unique continuation property for (1.1) fails for general L^p , $p < n$, potentials as demonstrated in Example 2.5, the following theorem shows that if the potential is in L^2 then the weak continuation property holds. Recall that weak unique continuation for a differential (in)equality is the property that every solution that vanishes in an open subset vanishes identically.

Theorem 2.7. *Let Ω be a domain in \mathbb{R}^n , $n \geq 2$, and let $V \in L^2_{loc}(\Omega)$. Suppose $u = (u_1, \dots, u_M) : \Omega \rightarrow \mathbb{R}^M$ with $u \in H^1_{loc}(\Omega)$ and satisfies $|\nabla u| \leq V|u|$ on Ω . If u vanishes in an open subset of Ω , then u vanishes identically.*

Proof. The $n = 2$ case is a direct consequence of Theorem 1.1, since the (strong) unique continuation implies the weak unique continuation property. We shall show below when $n = 3$, for any two domains D_1, D_2 in \mathbb{R}^2 with $D_1 \subseteq D_2$, and $s > 0$, if u satisfies (1.1) on the product domain $D_2 \times (-s, s)$ and $u = 0$ on $D_1 \times (-s, s)$, then $u = 0$ on $D_2 \times (-s, s)$. If so, then $u \equiv 0$ with a standard propagation argument. The proof for $n \geq 3$ cases follows from an induction.

Since $V \in L^2_{loc}(D_2 \times (-s, s))$, by Fubini's theorem, for almost every $x_3 \in (-s, s)$, $V(\cdot, x_3) \in L^2_{loc}(D_2)$, and similarly $u(\cdot, x_3) \in H^1_{loc}(D_2)$. Restricting (2.1) at each such $x_3 = c_3 \in (-s, s)$, we have $v = u(\cdot, c_3)$ satisfies

$$|\nabla v| = \left(\sum_{k=1}^M |\partial_{x_1} u_k(\cdot, c_3)|^2 + |\partial_{x_2} u_k(\cdot, c_3)|^2 \right)^{\frac{1}{2}} \leq |\nabla u(\cdot, c_3)| \leq V(\cdot, c_3)|u(\cdot, c_3)| = V(\cdot, c_3)|v|$$

on D_2 and $v = 0$ on D_1 . Applying the $n = 2$ case we have $v = 0$ on D_2 . Thus $u = 0$ on $D_2 \times (-s, s)$. \blacksquare

3. UNIQUENESS FOR LIPSCHITZ FUNCTIONS

In this section, we focus on locally Lipschitz functions whose definition is given below.

Definition 3.1. A function $u : \Omega \rightarrow \mathbb{R}$ is said to be locally Lipschitz on Ω means that for any point $p \in \Omega$, there is some neighborhood $p \in U_p \subseteq \Omega$ and some constant C_p so that for all $x, y \in U_p$, $|u(y) - u(x)| < C_p|y - x|$. The function u is Lipschitz on Ω means that there exists a constant C such that for all $x, y \in \Omega$, $|u(y) - u(x)| < C|y - x|$.

According to Rademacher's Theorem, if u is locally Lipschitz on Ω , then ∇u is defined a.e. on Ω . See, for instance, [E, pp. 296]. Moreover,

Proposition 3.2. [E, pp. 294] *Let Ω be a domain in \mathbb{R}^n . Then u is locally Lipschitz on Ω if and only if $u \in W^{1,\infty}_{loc}(\Omega)$. \blacksquare*

Following the convention of [E], even for $u \in W^{1,\infty}_{loc}(\Omega)$ defined a.e. in Ω or with some measure zero set of discontinuities, there is a unique continuous function agreeing with u a.e., which we will also denote u .

To prove Theorem 1.3, we begin with a uniqueness property of Lipschitz functions in one real variable on an interval, making use of the following fundamental theorem of calculus for Lipschitz functions.

Proposition 3.3. [R, Theorem 7.20, Fundamental Theorem of Calculus] *If $u : [0, 1] \rightarrow \mathbb{R}$ is Lipschitz on $[0, 1]$, then for any $0 \leq a < b \leq 1$,*

$$u(b) - u(a) = \int_a^b u'(t) dt.$$

■

Lemma 3.4. *Let $\varphi : [0, 1] \rightarrow \mathbb{R}$ be Lipschitz on $[0, 1]$, with $\varphi(0) = 0$. If there exist $p \geq 1$ and a non-negative function $\lambda : [0, 1] \rightarrow \mathbb{R}$ with $\lambda \in L^p([0, 1])$ such that for a.e. $x \in (0, 1)$,*

$$|\varphi'(x)| \leq \lambda(x) |\varphi(x)| x^{\frac{1-p}{p}}, \quad (3.1)$$

then $\varphi \equiv 0$ in $[0, 1]$.

Proof. We note first that we can assume without loss of generality that λ is non-vanishing. Indeed, if that is not the case, then we can just replace λ with $1 + \lambda \in L^p([0, 1])$ and (3.1) still holds.

Let $\delta = \sup\{d \in [0, 1] \mid \varphi \equiv 0 \text{ in } [0, d]\}$. By continuity, $\varphi(\delta) = 0$, and by Proposition 3.3 (which uses the Lipschitz hypothesis), for all $x \in (0, 1]$,

$$|\varphi(x)| = |\varphi(x) - \varphi(\delta)| = \left| \int_\delta^x \varphi'(t) dt \right| \leq \int_\delta^x |\varphi'(t)| dt. \quad (3.2)$$

The existence of the RHS integral is from Proposition 3.2, with $L^\infty([\delta, x]) \subseteq L^p([\delta, x]) \subseteq L^1([\delta, x])$.

For $p > 1$, let q be the conjugate exponent so that $\frac{1}{p} + \frac{1}{q} = 1$. By Hölder's inequality, we have

$$\int_\delta^x |\varphi'(t)| dt \leq \left(\int_\delta^x |\varphi'(t)|^p dt \right)^{\frac{1}{p}} \left(\int_\delta^x 1^q dt \right)^{\frac{1}{q}} \leq \left(\int_\delta^x |\varphi'(t)|^p dt \right)^{\frac{1}{p}} x^{\frac{p-1}{p}}. \quad (3.3)$$

It follows from (3.2) and (3.3) that for $p \geq 1$,

$$|\varphi(x)|^p \leq x^{p-1} \int_\delta^x |\varphi'(t)|^p dt. \quad (3.4)$$

We multiply both sides of (3.4) by the function $x^{1-p} \lambda^p(x)$, to obtain:

$$\lambda^p(x) |\varphi(x)|^p x^{1-p} \leq \lambda^p(x) \int_\delta^x |\varphi'(t)|^p dt. \quad (3.5)$$

Suppose toward a contradiction that $\delta < 1$, and let $s \in (\delta, 1)$. Integrating in the variable x on both sides of (3.5) gives

$$\int_{\delta}^s \lambda^p(x) |\varphi(x)|^p x^{1-p} dx \leq \int_{\delta}^s \left(\lambda^p(x) \int_{\delta}^x |\varphi'(t)|^p dt \right) dx. \quad (3.6)$$

Note that the Lipschitz property of φ on $[0, 1]$ and $\varphi(0) = 0$ imply there is some constant C so that $|\varphi(x)| \leq C|x|$, so $|\varphi(x)|^p x^{1-p}$ is continuous and bounded as a function of x on $(0, 1]$. Then the hypothesis $\lambda \in L^p([0, 1])$ applies, so that both the LHS and RHS integrals in (3.6) exist. The inequality (3.6) then implies, first using $x \leq s$, and then the hypothesis (3.1):

$$\begin{aligned} \int_{\delta}^s \lambda^p(x) |\varphi(x)|^p x^{1-p} dx &\leq \left(\int_{\delta}^s \lambda^p(x) dx \right) \left(\int_{\delta}^s |\varphi'(x)|^p dx \right) \\ &\leq \left(\int_{\delta}^s \lambda^p(x) dx \right) \left(\int_{\delta}^s \lambda^p(x) |\varphi(x)|^p x^{1-p} dx \right). \end{aligned} \quad (3.7)$$

By the construction of δ as the supremum of a set where $\varphi(x) \equiv 0$, we can find a sequence of points $s_j \in (\delta, 1)$ so that s_j is decreasing, $\lim_{j \rightarrow \infty} s_j = \delta$, and $\varphi(s_j) \neq 0$. By the continuity of $|\varphi(x)|^p x^{1-p}$ and the property that $\lambda^p \geq 1$, the integrand $\lambda^p(x) |\varphi(x)|^p x^{1-p}$ is strictly positive in some neighborhood of s_j . So, for all $j = 1, 2, 3, \dots$,

$$\int_{\delta}^{s_j} \lambda^p(x) |\varphi(x)|^p x^{1-p} dx > 0.$$

The inequality (3.7) then yields, for all j ,

$$1 \leq \int_{\delta}^{s_j} \lambda^p(x) dx. \quad (3.8)$$

Since $\lambda \in L^p(0, 1)$, letting $s_j \rightarrow \delta$ in (3.8) leads to a contradiction. \blacksquare

Recalling Rademacher's theorem that a Lipschitz function is differentiable almost everywhere, the following simple, but useful, Lemma gives a set of points where the square of a Lipschitz function is known to be differentiable.

Lemma 3.5. *Let u be a locally Lipschitz function on an open set $\Omega \subseteq \mathbb{R}^n$. Then $g = u^2$ is also locally Lipschitz on Ω . Moreover, g is differentiable wherever u vanishes and in fact $\nabla g(x) = 0$ there.*

Proof. The locally Lipschitz property of g follows from the well-known fact that the product of locally Lipschitz functions is locally Lipschitz; this is easily checked as an elementary consequence of Definition 3.1. The second claim is also elementary; let $x_0 \in \Omega$ be such that $u(x_0) = 0$. The properties that $g = u^2$ is differentiable at x_0 and $\nabla g(x_0) = 0$ follow from the definition of differentiability,

$$\lim_{x \rightarrow x_0} \frac{g(x) - g(x_0) - 0 \cdot (x - x_0)}{|x - x_0|} = \lim_{x \rightarrow x_0} \frac{u^2(x)}{|x - x_0|} = 0,$$

where we have used the Lipschitz property of $u(x) = u(x) - u(x_0) = O(|x - x_0|)$ in a neighborhood of x_0 . \blacksquare

Lemma 3.6. *For $n \geq 2$, let A be a set of measure zero in the unit ball in \mathbb{R}^n . Then for almost all points ω in the unit sphere S^{n-1} , the set of intersection of A with the radius segment $\{r\omega : 0 \leq r \leq 1\}$ is of measure zero in the line measure.*

Proof. Let $|K|$ denote the d -dimensional measure of a measurable set $K \subseteq \mathbb{R}^d$, and let $\chi_A : \mathbb{R}^n \rightarrow \mathbb{R}$ be the characteristic function of the set A . We have

$$0 = |A| = \int_{|x| < 1} \chi_A(x) dv = \int_{S^{n-1}} \int_0^1 \chi_A(r\omega) r^{n-1} dr d\omega.$$

By Fubini's theorem, we conclude that for a.e. $\omega \in S^{n-1}$, $\int_0^1 \chi_A(r\omega) r^{n-1} dr = 0$, which is the desired result: $|A \cap \{r\omega\}| = 0$. \blacksquare

Given a locally Lipschitz function u on Ω , denote by Z_u be the zero set of u in Ω , that is, $Z_u = \{x \in \Omega \mid u(x) = 0\}$. Theorem 1.3 will be a consequence of the following general result concerning Lipschitz functions.

Theorem 3.7. *Let Ω be a domain in \mathbb{R}^n , $n \geq 2$, and u be a locally Lipschitz function on Ω . If the zero set Z_u of u is neither \emptyset nor Ω , then*

$$\int_{\Omega \setminus Z_u} \left| \frac{\nabla u(x)}{u(x)} \right|^n dv = \infty. \quad (3.9)$$

Proof. First we make the observation that in order to prove the theorem it suffices to prove it for $g = u^2$. In fact, since

$$\int_{\Omega \setminus Z_u} \left| \frac{\nabla g(x)}{g(x)} \right|^n dv = 2^n \int_{\Omega \setminus Z_u} \left| \frac{\nabla u(x)}{u(x)} \right|^n dv,$$

if the conclusion (3.9) is true for $g = u^2$, then it is also true for u . Hence we only need to prove (3.9) for a function that is the square of a locally Lipschitz function. By Lemma 3.5, the gradient of g is 0 at every point where g is 0 (the same set Z_u where u is 0), and g also satisfies the locally Lipschitz assumption. For the rest of the proof we assume (by replacing u with g) that $\nabla u(x) = 0$ wherever $u(x) = 0$. Let

$$V(x) = \begin{cases} \left| \frac{\nabla u(x)}{u(x)} \right| & x \in \Omega \setminus Z_u \\ 0 & x \in Z_u. \end{cases} \quad (3.10)$$

Note that V is a measurable function in Ω . The zero set Z_u is closed in Ω , and by the assumptions that $Z_u \neq \emptyset$, $Z_u \neq \Omega$, and Ω is connected, there is some boundary point $x_0 \in \partial Z_u \subseteq Z_u \subseteq \Omega$, and a ball $B_{r_0}(x_0)$ such that the closure $\overline{B_{r_0}(x_0)} \subseteq \Omega$, and u is Lipschitz on $\overline{B_{r_0}(x_0)}$. We can assume, after a translation and scaling, that x_0 is the origin and the

radius r_0 is equal to 1. Because $B_1 \setminus Z_u$ is open and non-empty, there is some $B_{r_1}(x_1) \subseteq B_1$ where u is nonvanishing.

Now suppose toward a contradiction that (3.9) is false:

$$\int_{\Omega} V^n(x) dv = \int_{\Omega \setminus Z_u} \left| \frac{\nabla u(x)}{u(x)} \right|^n dv < \infty, \quad (3.11)$$

and therefore $V \in L^n(\Omega)$. Hence in polar coordinates,

$$\int_{B_1} V^n(x) dv = \int_{S^{n-1}} \int_0^1 V^n(r\omega) r^{n-1} dr d\omega. \quad (3.12)$$

Since the integral (3.12) is finite, Fubini's theorem implies that for a.e. $\omega \in S^{n-1}$ we have

$$\int_0^1 V^n(r\omega) r^{n-1} dr < \infty. \quad (3.13)$$

From Rademacher's theorem, let $A \subseteq \Omega$ be the set of measure zero where $\nabla u(x)$ does not exist at x . Choose $\omega_0 \in S^{n-1}$ such that (3.13) holds, that is, $V(r\omega_0)r^{\frac{n-1}{n}} \in L^n([0, 1])$ and at the same time, by Lemma 3.6 the same $\omega_0 \in S^{n-1}$ can be chosen such that $\nabla u(x)$ exists a.e. on the radius segment $\{r\omega_0\}$. Define φ , for $t \in [0, 1]$, by

$$\varphi(t) = u(t\omega_0).$$

It is evident that $\varphi(t)$ is Lipschitz on $[0, 1]$ from the Lipschitz property of u . Then applying the chain rule at points t such that u is differentiable at $t\omega_0$, we have

$$\varphi'(t) = \nabla u(t\omega_0) \cdot \omega_0,$$

which implies, for a.e. $t \in [0, 1]$,

$$|\varphi'(t)| \leq |\nabla u(t\omega_0)|. \quad (3.14)$$

By the definition of V , we have

$$|\varphi'(t)| \leq V(t\omega_0)|u(t\omega_0)| = V(t\omega_0)|\varphi(t)| \quad \text{for } u(t\omega_0) \neq 0. \quad (3.15)$$

However, when $u(t\omega_0) = 0$, we have, by the observation at the beginning of the proof, $\nabla u(r\omega_0) = 0$ and therefore $\varphi'(t) = 0$. Hence we have shown that

$$|\varphi'(t)| \leq V(t\omega_0)|\varphi(t)| = V(t\omega_0)t^{\frac{n-1}{n}}|\varphi(t)|t^{-\frac{n-1}{n}}$$

holds for a.e. $t \in [0, 1]$. By Lemma 3.4, with $\lambda(t) = V(t\omega_0)t^{\frac{n-1}{n}}$ and $p = n$, $\varphi(t) \equiv 0$. So $u \equiv 0$ on all the radius segments $\{t\omega_0\}$ for a.e. $\omega_0 \in S^{n-1}$, but this contradicts the fact that u has no zeros in the ball $B_{r_1}(x_1)$. \blacksquare

Remark 3.8. The proof of Theorem 3.7 actually leads to the following stronger conclusion: under the same assumption as in Theorem 3.7, on every neighborhood $U \subseteq \Omega$ of a point $a \in \Omega \cap \partial Z_u$, one has

$$\int_{U \setminus Z_u} \left| \frac{\nabla u(x)}{u(x)} \right|^n dv = \infty.$$

Proof of Theorem 1.3. For the one-dimensional case $n = 1$, Ω is an open interval (a, b) and Lemma 3.4 can be used directly. For any $s \in (x_0, b)$, let $\varphi(x) = u((s - x_0)x + x_0)$ so that $\varphi(0) = u(x_0) = 0$, $\varphi(1) = u(s)$, and φ is Lipschitz on $[0, 1]$. Lemma 3.4 applies to φ with $p = 1$ and $\lambda(x) = V((s - x_0)x + x_0) \cdot |s - x_0| \in L^1([0, 1])$, to show $\varphi(1) = 0 = u(s)$. Similarly, $u(t) = 0$ for any $a < t < x_0$.

For $n \geq 2$, suppose u and V satisfy $|\nabla u| \leq V|u|$ a.e. on Ω with $V \in L^n_{loc}(\Omega)$. Let $B = B_r(x_0)$ be a nonempty open ball such that $\overline{B} \subset \Omega$. Suppose toward a contradiction that $B \not\subseteq Z_u$. Then u is locally Lipschitz on B , and the zero set $Z_u \cap B$ of u in B is neither \emptyset (since $x_0 \in Z_u \cap B$) nor B . However,

$$\int_{B \setminus Z_u} \left| \frac{\nabla u(x)}{u(x)} \right|^n dv \leq \int_{B \setminus Z_u} |V(x)|^n dv < \infty,$$

contradicting Theorem 3.7. We can conclude $B \subseteq Z_u$. Thus Z_u is both open and closed in the connected set Ω and $u \equiv 0$. \blacksquare

Remark 3.9. The Lipschitz condition cannot just be dropped in Theorem 1.3 when $n \geq 2$. Indeed, Example 2.4 gives a nontrivial function u_0 that is locally Lipschitz on $B_{\frac{1}{2}} \setminus \{0\}$, continuous on $B_{\frac{1}{2}}$ with $u_0(0) = 0$, and solves $|\nabla u| = V|u|$ on $B_{\frac{1}{2}}$ for some $V \in L^n(B_{\frac{1}{2}})$. This indicates that the zero set of solutions fails to propagate at a non-Lipschitz point in general.

On the other hand, the hypothesis of Theorem 1.3 can be weakened as in the following Corollary without contradicting Remark 3.9 — if u is a continuous k^{th} root of a locally Lipschitz function, then the uniqueness still holds.

Corollary 3.10. *Let Ω be a domain in \mathbb{R}^n and $u : \Omega \rightarrow \mathbb{R}$ be continuous on Ω with the zero set $Z_u \subseteq \Omega$. If there is some integer $k \geq 1$ so that $v(x) = (u(x))^k$ is locally Lipschitz on Ω , then ∇u exists a.e. in $\Omega \setminus Z_u$. Further, if there is some $V \in L^n_{loc}(\Omega)$, so that ∇u satisfies*

$$|\nabla u| \leq V|u| \quad \text{a.e. on } \Omega \setminus Z_u,$$

and $Z_u \neq \emptyset$, then $u \equiv 0$.

Proof. On the open set where $u(x) > 0$, the partial derivatives of v exist a.e. and at each point where ∇v exists, by the chain rule, the partial derivatives of $u(x) = (v(x))^{1/k}$ exist, with $\nabla u(x) = \frac{1}{k}(v(x))^{\frac{1}{k}-1} \nabla v(x)$. Similarly, on the open set where $u(x) < 0$, the partial derivatives of $u = -((-1)^k v)^{1/k}$ exist a.e., establishing the first claim.

Consider $g(x) = (v(x))^2 = (u(x))^{2k}$. By Lemma 3.5, g is locally Lipschitz on Ω and at every point where $v(x) = 0$, which is the same set as Z_u , the partial derivatives of g exist with $\nabla g = 0$. At every point where $u(x) \neq 0$, if ∇u exists, then ∇g also exists and is equal to $2ku^{2k-1} \nabla u$. So, at every point in Ω except for a set of measure zero contained in $\Omega \setminus Z_u$, ∇g exists and satisfies:

$$|\nabla g| = 2k|u|^{2k-1} |\nabla u| \leq 2kV|u|^{2k} = 2kV|g|.$$

Applying Theorem 1.3 and the assumption $Z_u \neq \emptyset$, we have $g \equiv 0$, and thus $u \equiv 0$. \blacksquare

We conclude the section with an application of Theorem 1.3 to a uniqueness problem for a nonlinear system of differential equations.

Corollary 3.11. *For a domain $\Omega \subseteq \mathbb{R}^n$, $u : \Omega \rightarrow \mathbb{R}$, $x_0 \in \Omega$, $y_0 \in \mathbb{R}$, let $f : \mathbb{R} \rightarrow \mathbb{R}^n$ be Lipschitz on \mathbb{R} . Then there exists at most one Lipschitz solution to $\nabla u = f(u)$ on Ω with $u(x_0) = y_0$.*

Proof. Suppose there exists a pair of Lipschitz solutions u_1, u_2 to $\nabla u = f(u)$ on Ω with the same initial condition $u_1(x_0) = u_2(x_0) = y_0$. Then $w = u_1 - u_2$ is Lipschitz on Ω , $w(x_0) = 0$, and w satisfies

$$|\nabla w| = |\nabla u_1 - \nabla u_2| = |f(u_1) - f(u_2)| \leq C|u_1 - u_2| = C|w| \quad \text{on } \Omega,$$

where $C > 0$ is the Lipschitz constant for f . By Theorem 1.3, we have $w \equiv 0$. ■

4. FURTHER APPLICATIONS

Corollary 4.1. *Given any nonempty closed set $A \subsetneq \mathbb{R}^n$, there exists a smooth function $F : \mathbb{R}^n \setminus A \rightarrow \mathbb{R}$ so that for any point $a \in \partial A$, F cannot be extended to an L^n integrable function over any neighborhood of a .*

Proof. First, by a well-known theorem of Whitney ([W]), there exists a smooth function $h : \mathbb{R}^n \rightarrow \mathbb{R}$ whose zero set is exactly A . Its square, $u(x) = (h(x))^2$, is also smooth, has zero set exactly A , and satisfies $\nabla u = 0$ at every point of A by Lemma 3.5. The quotient $\frac{|\nabla u|}{|u|}$ is the claimed smooth function F on the open set $\mathbb{R}^n \setminus A$. If there were some ball $B = B_r(a)$ and a function $V \in L^n(B)$ which agrees with $\frac{|\nabla u|}{|u|}$ on $B \setminus A$, then u and V would satisfy (1.1) from Theorem 1.3 at every point of $B \setminus A$ by construction of V , and at every point of $B \cap A$, where $|\nabla u| = 0$. By Theorem 1.3, $u \equiv 0$ on B , contradicting the assumption that a is a boundary point of the zero set. ■

Theorem 4.2. *Let Ω be an open set in \mathbb{R}^n , and u be a Lipschitz function on Ω . Then $\log |u(x) - u(a)| \notin W_{loc}^{1,n}(\Omega \setminus Z_{u-u(a)})$ for every $a \in \Omega$. In particular, if $\log |u(x)| \in W_{loc}^{1,n}(\Omega \setminus Z_u)$, then u is nowhere zero on Ω . If, in addition, Ω has Lipschitz boundary, then the above results also hold true with Ω replaced by $\bar{\Omega}$.*

Proof. Let $v = u - u(a)$ on Ω . Then v is Lipschitz on Ω with $v(a) = 0$. If $\log |v| \notin L_{loc}^n(\Omega)$, then we are done. If $\log |v| \in L_{loc}^n(\Omega)$, then one further computes

$$|\nabla \log |v|| = \frac{|\nabla v|}{|v|}$$

wherever $v \neq 0$. If a is an interior point of $Z_v = \{x \in \Omega \mid v(x) = 0\}$, the zero set of v , then the theorem is trivially true. If $a \in \partial Z_v \cap \Omega$, then one can apply Remark 3.8 to conclude $\nabla \log |v| \notin L_{loc}^n(\Omega \setminus Z_v)$.

In the case when $a \in \partial\Omega$ and Ω has Lipschitz boundary, if $v(b) = 0$ for some $b \in \Omega$, then it is reduced to the $a \in \Omega$ case. Thus we assume $a \in \partial\Omega$ and $v(x) \neq 0$ for all $x \in \Omega$. In

particular, this means there exists a cone $S_a \subseteq \Omega$ centered at a (which exists since Ω has Lipschitz boundary) such that $v \neq 0$ on S_a . Making use of a similar argument as in the proof of Theorem 3.7, with B_1 replaced by S_a , one can obtain $\nabla \log |v| \notin L^n_{loc}(\Omega)$. ■

A natural way to view Theorem 4.2 is as follows. Denote by $Lip(\Omega)$ the set of all Lipschitz functions on Ω . Theorem 4.2 implies that for each $a \in \Omega$,

$$T_a(Lip(\Omega)) \cap W^{1,n}(\Omega) = \emptyset,$$

where T_a is a (non-linear) map on $Lip(\Omega)$ defined by $T_a(u) = \log |u - u(a)|$, $u \in Lip(\Omega)$. The following Corollary is a direct consequence of Theorem 4.2 as well.

Corollary 4.3. *Let Ω be an open set in \mathbb{R}^n and $u : \Omega \rightarrow \mathbb{R}$ be locally Lipschitz on Ω . If the zero set Z_u of u is neither empty nor Ω , and*

$$\int_{\Omega \setminus Z_u} |\nabla \log |u(x)||^p dv < \infty. \quad (4.1)$$

then $p < n$. ■

Proposition 4.4. *Let Ω be an open set in \mathbb{R}^n , $n \geq 2$, and $\phi : \Omega \rightarrow \mathbb{R}$ with $\phi \in W^{1,n}_{loc}(\Omega)$. The following statements hold for the exponential $e^{-|\phi|}$.*

- (1) $e^{-|\phi|}$ vanishes to finite order in the L^2 sense anywhere in Ω .
- (2) If $e^{-|\phi|}$ is locally Lipschitz on Ω , then $e^{-|\phi|}$ is nowhere zero on Ω . $e^{-|\phi|}$ is nowhere zero on $\overline{\Omega}$ if in addition Ω has Lipschitz boundary.

Proof. Since $\phi \in W^{1,n}_{loc}(\Omega)$, we have $|\phi| \in W^{1,n}_{loc}(\Omega)$ as well. The function $u = e^{-|\phi|}$ satisfies $|u| < 1$ and

$$|\nabla u| = |\nabla |\phi|| e^{-|\phi|} = |\nabla |\phi|| |u| \leq |\nabla |\phi|| \in L^n_{loc}(\Omega).$$

See for instance [E, pp. 308]. Hence $u \in W^{1,n}_{loc}(\Omega)$ and satisfies $|\nabla u| = V|u|$ with $V = |\nabla |\phi|| \in L^n_{loc}(\Omega)$. By Theorem 1.1, u cannot vanish to infinite order in the L^2 sense anywhere in Ω .

If $e^{-|\phi|}$ is also Lipschitz on Ω , and $e^{-|\phi|}$ is zero at $x_0 \in \Omega$, then $|\phi| = -\log |u - u(x_0)| \notin W^{1,n}_{loc}(\Omega)$ by Theorem 4.2. Contradiction! ■

Before proving Theorem 1.2, let us recall the **Moser-Trudinger inequality**: Let Ω be a bounded domain in \mathbb{R}^n with Lipschitz boundary, and $\alpha_n = nw_{n-1}^{\frac{1}{n-1}}$ where w_{n-1} is the surface area of the unit sphere in \mathbb{R}^n . There exists a positive constant C_{MT} depending only on n such that

$$\sup_{u \in W^{1,n}_0(\Omega), \|\nabla u\|_{L^n(\Omega)} \leq 1} \int_{\Omega} e^{\alpha_n |u(x)|^{\frac{n}{n-1}}} dv \leq C_{MT} |\Omega|.$$

Here $|\Omega|$ is the volume of Ω . We shall use the Moser-Trudinger inequality to prove that the exponential of $W^{1,n}$ functions is L^2 integrable.

Proof of Theorem 1.2: First we show that $e^\phi \in L_{loc}^2(\Omega)$. Pick r small enough such that $B_{2r}(x_0) \subseteq \Omega$. By Sobolev extension theorem, there exists an extension $\tilde{\phi} \in W_0^{1,n}(B_{2r}(x_0))$ of $\phi|_{B_r(x_0)}$ such that

$$a = \|\nabla \tilde{\phi}\|_{L^n(B_{2r}(x_0))} \leq C \|\phi\|_{W^{1,n}(B_r(x_0))}$$

for some constant C dependent only on r and n . In particular, $\tilde{\phi}_1 := a^{-1}\tilde{\phi} \in W_0^{1,n}(B_{2r}(x_0))$ and $\|\nabla \tilde{\phi}_1\|_{L^n(B_{2r}(x_0))} \leq 1$. Thus one applies the Moser-Trudinger inequality to obtain

$$\int_{B_{2r}(x_0)} e^{\alpha_n |\tilde{\phi}_1(x)|^{\frac{n}{n-1}}} dv \lesssim 1.$$

Noting that $2\tilde{\phi} < \alpha_n |\tilde{\phi}_1|^{\frac{n}{n-1}}$ when $|\tilde{\phi}| > 2^{n-1} a^n \alpha_n^{1-n}$, we further have

$$\int_{B_{2r}(x_0) \cap \{|\tilde{\phi}| > 2^{n-1} a^n \alpha_n^{1-n}\}} e^{2\tilde{\phi}(x)} dv \leq \int_{B_{2r}(x_0)} e^{\alpha_n |\tilde{\phi}_1(x)|^{\frac{n}{n-1}}} dv \lesssim 1.$$

The claim that $e^\phi \in L_{loc}^2(\Omega)$ is thus a consequence of the following inequality

$$\begin{aligned} \int_{B_r(x_0)} e^{2\phi(x)} dv &\leq \int_{B_{2r}(x_0) \cap \{|\tilde{\phi}| \leq 2^{n-1} a^n \alpha_n^{1-n}\}} e^{2\tilde{\phi}(x)} dv + \int_{B_{2r}(x_0) \cap \{|\tilde{\phi}| > 2^{n-1} a^n \alpha_n^{1-n}\}} e^{2\tilde{\phi}(x)} dv \\ &\lesssim e^{2^n a^n \alpha_n^{1-n}} r^n + 1. \end{aligned}$$

On the other hand, by Proposition 4.4 part (1), $e^{-|\phi|}$ vanishes to finite order in the L^2 sense at x_0 . Equivalently, there exists some $m_0 \geq 0$ such that

$$\overline{\lim}_{r \rightarrow 0} r^{-m_0} \int_{|x-x_0| < r} |e^{-|\phi(x)|}|^2 dv > 0.$$

Since $e^\phi \geq e^{-|\phi|} \geq 0$ and $e^\phi \in L_{loc}^2(\Omega)$, we further have

$$\overline{\lim}_{r \rightarrow 0} r^{-m_0} \int_{|x-x_0| < r} |e^{\phi(x)}|^2 dv \geq \overline{\lim}_{r \rightarrow 0} r^{-m_0} \int_{|x-x_0| < r} |e^{-|\phi(x)|}|^2 dv > 0.$$

Namely, e^ϕ vanishes to finite order in the L^2 sense at x_0 . ■

Corollary 4.5. *Let Ω be an open set in \mathbb{R}^n , $n \geq 2$. Suppose $\phi : \Omega \rightarrow \mathbb{R}$ with $\phi \in W_{loc}^{1,n}(\Omega)$. If e^ϕ is Lipschitz on Ω , then e^ϕ is nowhere zero on Ω .*

Proof. It is not hard to verify that for all $x_1, x_2 \in \Omega$,

$$|e^{-|\phi(x_2)|} - e^{-|\phi(x_1)|}| \leq |e^{\phi(x_1)} - e^{\phi(x_2)}|.$$

In particular, $e^{-|\phi|}$ is Lipschitz whenever e^ϕ is so. Applying Proposition 4.4 part (2), we have $e^{-|\phi|}$, and thus e^ϕ , is nowhere zero on Ω . ■

5. IN RELATION TO $\bar{\partial}$

On domains in \mathbb{C}^n , if the gradient operator ∇ is replaced by the $\bar{\partial}$ operator, then Theorem 1.3 fails, even for real analytic functions. In fact, there are real analytic functions that vanish to any given order at one point and satisfy $|\bar{\partial}u| \leq V|u|$ for some $V \in L^\infty$.

Example 5.1. Let f be a holomorphic function on $B_1 \subseteq \mathbb{C}^n$ that vanishes to order k at 0, $k \geq 1$. Letting $u(z) = (1 + \frac{z_1}{2})f(z)$, then u is real analytic on B_1 , vanishes to order k at 0 and satisfies $|\bar{\partial}u| \leq 4|u|$.

On the other hand, since $|\nabla u|^2 = |\partial u|^2 + |\bar{\partial}u|^2$ for a Lipschitz u , by Theorem 1.3 we have near any neighborhood U of a zero point in ∂Z_u of u ,

$$\int_U \frac{|\nabla u(z)|^2}{|u(z)|^2} dv = \int_U \frac{|\bar{\partial}u(z)|^2}{|u(z)|^2} + \frac{|\partial u(z)|^2}{|u(z)|^2} dv = \infty.$$

The following propositions discuss a finer property about the L^2 divergence of $\frac{\nabla u}{u}$ concerning the smooth extension of holomorphic functions beyond the boundary. In particular, they exhibit an intrinsic obstruction for holomorphic functions to be extended smoothly across the boundary. We note that for smooth functions, the flatness in the L^2 sense at a point is equivalent to the vanishing of all jets at that point.

Proposition 5.2. *Let Ω be a domain in \mathbb{C} and $z_0 \in \partial\Omega$. Let u be a nonconstant holomorphic function on Ω . If u can be extended smoothly across z_0 , still denoted by u , and $u(z_0) = 0$, then there exists a neighborhood U of z_0 such that one of the following holds.*

(1) *If u vanishes to finite order at z_0 , then*

$$\int_U \frac{|\bar{\partial}u(z)|^2}{|u(z)|^2} dv < \infty \quad \text{and} \quad \int_U \frac{|\partial u(z)|^2}{|u(z)|^2} dv = \infty. \quad (5.1)$$

(2) *If u vanishes to infinite order at z_0 , then*

$$\int_U \frac{|\bar{\partial}u(z)|^2}{|u(z)|^2} dv = \infty. \quad (5.2)$$

Proof. Without loss of generality let $z_0 = 0$. In (1), since u vanishes to finite order at 0 and is holomorphic on Ω , $u = cz^k + O(|z|^{k+1})$ near 0 for some constant $c \neq 0, k \in \mathbb{Z}^+$. With a direct computation we have

$$\frac{|\bar{\partial}u|}{|u|} = \frac{O(z^k)}{|cz^k + O(|z|^{k+1})|} = O(1) \quad \text{and} \quad \frac{|\partial u|}{|u|} = \frac{k}{|z|} + O(1), \quad (5.3)$$

from which (5.1) follows.

For (2), if not, then set $V = \frac{\bar{\partial}u}{u}$ where $u \neq 0$, and $V = 0$ otherwise on U , so that $V \in L^2(U)$ and $\bar{\partial}u = Vu$ on U . According to Theorem 2.3, since u is flat at z_0 , we have $u \equiv 0$ on U . In particular, $u = 0$ on the open set $U \cap \Omega$. By the holomorphic property of u on Ω , we further have $u \equiv 0$ on Ω . This contradicts the assumption that u is nonconstant on Ω . \blacksquare

The following two corollaries give alternative characterizations on the vanishing order of smooth extension of holomorphic functions across the boundary.

Corollary 5.3. *Let Ω be a domain in \mathbb{C} and $z_0 \in \partial\Omega$. Let u be a nonconstant holomorphic function on Ω , and extend smoothly across z_0 , still denoted by u , with $u(z_0) = 0$. Then the following statements are equivalent to each other.*

- (1) u vanishes to finite order at z_0 .
- (2) $\frac{|\bar{\partial}u|}{|u|} \in L^\infty$ near z_0 .
- (3) $\frac{|\bar{\partial}u|}{|u|} \in L^2$ near z_0 .

Corollary 5.4. *Let Ω be a domain in \mathbb{C} and $z_0 \in \partial\Omega$. Let u be a nonconstant holomorphic function on Ω , and extend smoothly across z_0 , still denoted by u , with $u(z_0) = 0$. Then the following statements are equivalent to each other.*

- (1) u vanishes to infinite order at z_0 .
- (2) $\frac{|\bar{\partial}u|}{|u|} \notin L^\infty$ near z_0 .
- (3) $\frac{|\bar{\partial}u|}{|u|} \notin L^2$ near z_0 .

Proof of Corollary 5.3 and 5.4: For Corollary 5.3, (2) \Rightarrow (3) is trivial. (3) \Leftrightarrow (1) is a direct consequence of Proposition 5.2. (1) \Rightarrow (2) follows from (5.3) in the proof of Proposition 5.2. Corollary 5.4 can be proved similarly. ■

Example 5.5. Let \mathbb{H}^+ be the upper half plane in \mathbb{C} . The function

$$u = \exp\left(\frac{1}{i\sqrt{iz}}\right), \quad \arg iz \in \left(\frac{\pi}{2}, \frac{3\pi}{2}\right),$$

is holomorphic on \mathbb{H}^+ and vanishes to infinite order at $z_0 = 0$. It allows for a smooth extension across 0. By Proposition 5.2 (2), every smooth extension of u on a neighborhood U of 0 should satisfy (5.2). Note that u cannot extend holomorphically across 0.

For every $k \geq 1$, the function $u = z^k$ is holomorphic on \mathbb{H}^+ and vanishes to finite order k at 0. By Proposition 5.2 (1), every smooth extension of u on a neighborhood U of 0 should satisfy (5.1). For a less trivial example towards Proposition 5.2 (1) without holomorphic extension across 0, one can consider $u = z^k + e^{\frac{1}{i\sqrt{iz}}}$ on \mathbb{H}^+ instead, and obtain (5.1) for every smooth extension of u across 0.

Proposition 5.6. *Let Ω be a domain in \mathbb{C}^n and $z_0 \in \partial\Omega$. Let u be a nonconstant holomorphic function on Ω . If u can be extended smoothly across z_0 , still denoted by u , and $u(z_0) = 0$, then there exists a neighborhood U of z_0 such that one of the following holds.*

- (1) *If u vanishes to finite order at z_0 , then there exists a complex line L passing through z_0 such that*

$$\int_{U \cap L} \frac{|\partial u(z)|^2}{|u(z)|^2} dv = \infty. \quad (5.4)$$

(2) If u vanishes to infinite order at z_0 , then for every complex line L passing through z_0 ,

$$\int_{U \cap L} \frac{|\bar{\partial}u(z)|^2}{|u(z)|^2} dv = \infty. \quad (5.5)$$

Proof. For simplicity let $z_0 = 0$ and $n = 2$. The higher dimensional cases can be proved similarly. If u vanishes to finite order at 0, then after a holomorphic change of coordinates, there exists some $k \in \mathbb{Z}^+$ such that

$$u = z_1^k + g_{k-1}(z_2)z_1^{k-1} + \cdots + g_0(z_2) + h(z)$$

near 0. Here for each $j = 0, \dots, k-1$, g_j is smooth on U , holomorphic on $\Omega \cap U$ and $g_j(0) = 0$, and h is a smooth function on U with $h = 0$ on $\Omega \cap U$. In particular, h is flat at 0. Thus on the complex line $L = \{(z_1, 0) \in \mathbb{C}^2\}$, we have $u|_{U \cap L} = z_1^k + h(z_1, 0)$ and so

$$\frac{|\partial_{z_1} u|}{|u|} = \frac{k}{|z_1|} + O(1).$$

Hence (5.4) holds.

Assume u vanishes to infinite order at 0 and there exists a complex line L through 0 such that $\frac{|\bar{\partial}u|}{|u|} \in L^2(U \cap L)$. Applying a holomorphic change of coordinates if necessary, one can always write $L = \{(z_1, 0) \in \mathbb{C}^2\}$. Then $v := u|_{U \cap L}$ vanishes to infinite order at 0 and

$$\frac{|\bar{\partial}_{z_1} u|}{|u|} \leq \frac{|\bar{\partial}u|}{|u|} \in L^2(U \cap L).$$

In particular, there exists some $W \in L^2(U \cap L)$ such that $\bar{\partial}_{z_1} v = Wv$ on $U \cap L$. By Theorem 2.3, we have $v \equiv 0$. Thus (5.5) holds. \blacksquare

Proposition 5.7. *Let Ω be a domain in \mathbb{C}^n and $z_0 \in \partial\Omega$. Let u be a function holomorphic on Ω and smooth on a neighborhood $U \subseteq \mathbb{C}^n$ of z_0 . Then one of the following mutually exclusive cases holds.*

(1) u is holomorphic on $\Omega \cup U$.

(2)

$$\int_{U \setminus Z_{\bar{\partial}u}} \frac{\sum_{j,k=1}^n |\bar{\partial}_{z_j z_k}^2 u(z)|^2}{\sum_{j=1}^n |\bar{\partial}_{z_j} u(z)|^2} dv = \infty, \quad (5.6)$$

where $Z_{\bar{\partial}u}$ is the zero set of the vector function $\bar{\partial}u$.

Proof. Suppose u is not holomorphic on $\Omega \cup U$ and (5.6) fails. Then $Z_{\bar{\partial}u} \cap U \neq U$ and the function

$$W = \begin{cases} \sqrt{\frac{\sum_{j,k=1}^n |\bar{\partial}_{z_j z_k}^2 u|^2}{\sum_{j=1}^n |\bar{\partial}_{z_j} u|^2}}, & \text{on } U \setminus Z_{\bar{\partial}u}; \\ 0, & \text{on } Z_{\bar{\partial}u} \end{cases}$$

belongs to $L^2_{loc}(U)$. Let $v = (\bar{\partial}_{z_1} u, \dots, \bar{\partial}_{z_n} u)$. Then $v : U \rightarrow \mathbb{C}^n$ satisfies $|\bar{\partial}v| = W|v|$ on U and vanishes on the nonempty open set $U \cap \Omega$. According to the weak unique continuation

property in [PZ, Theorem 1.2], we have $v \equiv 0$ on U , contradicting the assumption that $Z_{\bar{\partial}_u} \cap U \neq U$. \blacksquare

We point out that in the case when Ω is pseudoconvex with smooth boundary, there always exists a function which is holomorphic on Ω and smooth on $\bar{\Omega}$, but does not extend holomorphically across a boundary point z_0 . Thus for every smooth extension of this function, part (2) of Proposition 5.7 always occurs.

Remark 5.8. While all the propositions and corollaries in this section are formulated for holomorphic functions with smooth extension across a boundary point, the same reasoning and conclusions can be extended without effort to more general settings, including formally holomorphic functions – smooth functions where the Taylor expansion at that point does not contain \bar{z} terms. See [FP] for more discussion on formally holomorphic functions.

Acknowledgments. Part of this work by the second named author was conducted while on sabbatical leave visiting Huaqiao university in China in Spring 2024. He thanks Jianfei Wang for his invitation, and the host institution for its hospitality and excellent research environment.

We also acknowledge the helpful comments of the reviewers, whose suggestions improved the writing and results of Sections 3 and 4.

REFERENCES

- [E] L. EVANS, *Partial Differential Equations*, Second edition. Graduate Studies in Mathematics **19**. American Mathematical Society, 2010.
- [FP] M. FASSINA and Y. PAN, *A local obstruction for elliptic operators with real analytic coefficients on flat germs*, Ann. Fac. Sci. Toulouse Math. Série 6 (5) **31** (2022), 1343–1363.
- [PW] Y. PAN and M. WANG, *Uniqueness for n -th order differential systems with strong singularities*, Electron. J. Differential Equations (2010), No. 172, 9 pp.
- [PZ] Y. PAN and Y. ZHANG, *Unique continuation for $\bar{\partial}$ with square-integrable potentials*, New York Journal of Mathematics **29** (2023), 402–416.
- [R] W. RUDIN, *Real and Complex Analysis*, Third edition. McGraw-Hill, 1987.
- [W] H. WHITNEY, *Analytic extensions of differentiable functions defined in closed sets*, Trans. Amer. Math. Soc. (1) **36** (1934), 63–89.

DEPARTMENT OF MATHEMATICAL SCIENCES, PURDUE UNIVERSITY FORT WAYNE, 2101 EAST COLISEUM BOULEVARD, FORT WAYNE, IN 46805, USA

Email address: coffmana@pfw.edu, pan@pfw.edu, zhangyu@pfw.edu