

Unique continuation for $\bar{\partial}$ with square-integrable potentials

Yifei Pan and Yuan Zhang

ABSTRACT. In this paper, we investigate the unique continuation property for the inequality $|\bar{\partial}u| \leq V|u|$, where u is a vector-valued function from a domain in \mathbb{C}^n to \mathbb{C}^N , and the potential $V \in L^2$. We show that the strong unique continuation property holds when $n = 1$, and the weak unique continuation property holds when $n \geq 2$. In both cases, the L^2 integrability condition on the potential is optimal.

1. Introduction

Let $u = (u_1, \dots, u_N)$ be a vector-valued function from a domain $\Omega \subset \mathbb{C}^n$ into \mathbb{C}^N . We say a differential equation or inequality satisfies the strong unique continuation property, if every solution that vanishes to infinite order at a point $z_0 \in \Omega$ vanishes identically. Here a square-integrable function u is said to vanish to infinite order (or flat) at $z_0 \in \Omega$ if for all $m \geq 0$,

$$(1.1) \quad \lim_{r \rightarrow 0} r^{-m} \int_{|z-z_0| < r} |u(z)|^2 dv_z = 0,$$

where dv_z is the Lebesgue measure element in \mathbb{C}^n with respect to the dummy variable z . A differential equation or inequality is said to satisfy the weak unique continuation property, if every solution that vanishes in an open subset vanishes identically.

While studying the boundary regularity and the uniqueness for CR-mappings of hypersurfaces in [2], Bell and Lempert had proved and applied the strong unique continuation property for $|\bar{\partial}u| \leq C|u|$ with C a constant. In this paper, we consider the unique continuation property for the following general type:

$$(1.2) \quad |\bar{\partial}u| \leq V|u| \quad \text{a.e.},$$

where the potential V is assumed to be locally square-integrable, i.e., $V \in L^2_{loc}(\Omega)$. This L^2 assumption on the potential is rather critical since the strong unique continuation property fails when $V \notin L^2$, as demonstrated by Example 2 in Section 3. Moreover, Mandache in [8] constructed an example (see Example 3) with an L^p potential, $0 < p < 2$, where even the weak unique continuation property fails.

2010 *Mathematics Subject Classification.* Primary 32W05; Secondary 35A23, 35A02.

Key words and phrases. Cauchy-Riemann equation, unique continuation, Hardy-Littlewood-Sobolev inequality.

Our first theorem below concerns the strong unique continuation property in the case when $n = 1$. Denote by $H_{loc}^k(\Omega)$ the standard (local) Sobolev space of functions whose weak derivatives up to order k are in $L_{loc}^2(\Omega)$.

Theorem 1.1. *Let Ω be a domain in \mathbb{C} . Suppose $u = (u_1, \dots, u_N) : \Omega \rightarrow \mathbb{C}^N$ with $u \in H_{loc}^1(\Omega)$ and satisfies $|\bar{\partial}u| \leq V|u|$ for some $V \in L_{loc}^2(\Omega)$. If u vanishes to infinite order at $z_0 \in \Omega$, then u vanishes identically.*

One can compare Theorem 1.1 with the following strong unique continuation property proved by Chanillo and Sawyer [3] for the differential inequality

$$(1.3) \quad |\Delta u| \leq V|\nabla u|$$

with the potential $V \in L^2$.

Theorem 1.2. [3] *Let Ω be a domain in \mathbb{C} . Suppose $u = (u_1, \dots, u_N) : \Omega \rightarrow \mathbb{R}^N$ with $u \in H_{loc}^2(\Omega)$ and satisfies $|\Delta u| \leq V|\nabla u|$ for some $V \in L_{loc}^2(\Omega)$. If u vanishes to infinite order at $z_0 \in \Omega$, then u vanishes identically.*

We note that the two inequalities (1.2)-(1.3) can easily be converted to each other, at least in the smooth category. However, in terms of the unique continuation property, it turns out Theorem 1.1 and Theorem 1.2 are no longer equivalent. In fact, one can obtain Theorem 1.2 from Theorem 1.1 by applying Theorem 1.1 to ∂u , as a consequence of the facts that $\Delta = 4\bar{\partial}\partial$ and $|\partial u(z)| = \frac{1}{2}|\nabla u(x, y)|$ (since u is real-valued in Theorem 1.2). To deduce Theorem 1.1 from Theorem 1.2, it would boil down to the possible existence of flat solutions to $\bar{\partial}$ given flat data. Surprisingly, there is an obstruction to such existence in the flat category. This phenomenon was discovered by Liu and the two authors in [7] which constructed a smooth function f that is flat at $0 \in \mathbb{C}$, yet $\bar{\partial}u = f$ has no solutions that are flat at 0. The same phenomenon was later investigated by Fassina and the first author in [5] for general elliptic operators with real analytic coefficients.

While Theorem 1.1 and Theorem 1.2 are concerned with local vector-valued solutions, there are earlier results about global and/or scalar (namely, $N = 1$) solutions to (1.2) and (1.3) on \mathbb{C} with $L^2(\mathbb{C})$ potentials. See, for instance, [6] by Kenig and Wang and [12] by Seo.

A well-known example of Wolff shows that the strong unique continuation for (1.3) no longer holds when the real dimension of the source domain is larger than 4. More precisely, Wolff constructed in [14] a smooth real-valued function on \mathbb{R}^d , $d \geq 5$, which vanishes to infinite order at the origin and satisfies (1.3) with $V \in L^d(\mathbb{R}^d)$.

As an immediate application of Theorem 1.1, we obtain the following weak unique continuation property for (1.2) with an L^2 potential when $n \geq 2$. Here given $u \in H_{loc}^1(\Omega)$, $\bar{\partial}u$ is understood as a $(0, 1)$ form with distribution components.

Theorem 1.3. *Let Ω be a domain in \mathbb{C}^n . Suppose $u = (u_1, \dots, u_N) : \Omega \rightarrow \mathbb{C}^N$ with $u \in H_{loc}^1(\Omega)$ and satisfies $|\bar{\partial}u| \leq V|u|$ for some $V \in L_{loc}^2(\Omega)$. If u vanishes in an open subset of Ω , then u vanishes identically.*

In contrast to the classical strong unique continuation property results for Laplacian, the integrability assumption on the potential in Theorem 1.3 is independent of the dimension n of the source domain. On the other hand, the weak unique continuation property fails for (1.2) with L^p potentials, $0 < p < 2$, as shown by a two-dimensional Mandache-type example (see Example 5 in Section 4). This shows the integrability assumption in Theorem 1.3 on the potential is optimal for the weak unique continuation property.

Theorem 1.3 can be applied to obtain the uniqueness of solutions as follows. Denote by $\mathbb{C}_{(0,1)}^N$ the space of N -vectors each of whose components is a $(0,1)$ form.

Corollary 1.4. *Let Ω be a domain in \mathbb{C}^n , $h = (h_1, \dots, h_N)^T : \Omega \rightarrow \mathbb{C}_{(0,1)}^N$, and $A = (A_{jk})_{1 \leq j, k \leq N} : \Omega \rightarrow \mathbb{C}_{(0,1)}^{N^2}$ with $A \in L_{loc}^2(\Omega)$. Suppose $f, g : \Omega \rightarrow \mathbb{C}^N$ with $f, g \in H_{loc}^1(\Omega)$ and are solutions to $\bar{\partial}u = Au + h$. If $f = g$ in an open subset of Ω , then $f \equiv g$ on Ω .*

2. Weighted Hardy-Littlewood-Sobolev inequality

Recall that given $f \in L^2(\mathbb{C})$, the Hardy-Littlewood Maximal function of f , denoted by Mf , is given by

$$Mf(z) := \sup \frac{1}{|D|} \int_D |f(\zeta)| dv_\zeta, \quad z \in \mathbb{C},$$

where the supremum is taken over all disks D in \mathbb{C} containing z . The Riesz fractional integral of f of order $\alpha \in (0, 2)$ is

$$I_\alpha f(z) := \int_{\mathbb{C}} \frac{f(\zeta)}{|\zeta - z|^{2-\alpha}} dv_\zeta, \quad z \in \mathbb{C}.$$

Definition 2.1. *Given a weight function $V(> 0)$ on a domain $\Omega \subset \mathbb{C}$, the weighted $L^2(\Omega)$ space with respect to V , denoted by $L_V^2(\Omega)$, is the collection of all functions f on Ω such that*

$$\|f\|_{L_V^2(\Omega)} := \left(\int_\Omega |f(z)|^2 V(z) dv_z \right)^{\frac{1}{2}} < \infty.$$

A more general but rather technical variant of the following theorem can be found in [4]. We shall prove the boundedness of

$$I_1 f = \int_{\mathbb{C}} \frac{f(\zeta)}{|\zeta - \cdot|} dv_\zeta$$

from $L_{V^{-1}}^2(\mathbb{C})$ space to $L_V^2(\mathbb{C})$ space with respect to a weight $V \in L^2(\mathbb{C})$ through a much simpler approach.

Theorem 2.2. *Let $V \in L^2(\mathbb{C})$. Then there exists a universal constant C_0 such that for any $f \in L^2_{V^{-1}}(\mathbb{C})$,*

$$\|I_1 f\|_{L^2_V(\mathbb{C})} \leq C_0 \|V\|_{L^2(\mathbb{C})} \|f\|_{L^2_{V^{-1}}(\mathbb{C})}.$$

Proof. Throughout the proof we use C to represent a universal constant, which may be different at different occurrences. We first show that for any $g \in L^2(\mathbb{C})$,

$$(2.1) \quad \|I_{\frac{1}{2}} g(z)\|_{L^2_V(\mathbb{C})} \leq C \|V\|_{L^2(\mathbb{C})}^{\frac{1}{2}} \|g\|_{L^2(\mathbb{C})}.$$

Without loss of generality, we assume $g \geq 0$.

For each $z \in \mathbb{C}$ with $\delta > 0$ to be chosen later, write

$$I_{\frac{1}{2}} g(z) = \int_{|\zeta-z|<\delta} + \int_{|\zeta-z|>\delta} \frac{g(\zeta)}{|\zeta-z|^{\frac{3}{2}}} dv_\zeta =: I + II.$$

By Hölder inequality,

$$\begin{aligned} II &\leq \left(\int_{|\zeta-z|>\delta} |g(\zeta)|^2 dv_\zeta \right)^{\frac{1}{2}} \left(\int_{|\zeta-z|>\delta} \frac{1}{|\zeta-z|^3} dv_\zeta \right)^{\frac{1}{2}} \\ &\leq C \|g\|_{L^2(\mathbb{C})} \left(\int_\delta^\infty \frac{1}{s^2} ds \right)^{\frac{1}{2}} = \frac{C}{\delta^{\frac{1}{2}}} \|g\|_{L^2(\mathbb{C})}. \end{aligned}$$

For I ,

$$\begin{aligned} I &= \sum_{k=1}^{\infty} \int_{\frac{\delta}{2^k} < |\zeta-z| < \frac{\delta}{2^{k-1}}} \frac{g(\zeta)}{|\zeta-z|^{\frac{3}{2}}} dv_\zeta \\ &\leq \sum_{k=1}^{\infty} \left(\frac{2^k}{\delta} \right)^{\frac{3}{2}} \int_{|\zeta-z| < \frac{\delta}{2^{k-1}}} g(\zeta) dv_\zeta \\ &= \sum_{k=1}^{\infty} \left(\frac{2^k}{\delta} \right)^{\frac{3}{2}} \left| D_{\frac{\delta}{2^{k-1}}} \right| \left(\frac{1}{\left| D_{\frac{\delta}{2^{k-1}}} \right|} \int_{|\zeta-z| < \frac{\delta}{2^{k-1}}} g(\zeta) dv_\zeta \right) \\ &\leq \sum_{k=1}^{\infty} 2^{-\frac{k}{2}+2} \pi \delta^{\frac{1}{2}} M g(z) \\ &= C \delta^{\frac{1}{2}} M g(z). \end{aligned}$$

Thus we have

$$I_{\frac{1}{2}} g(z) \leq C \left(\delta^{\frac{1}{2}} M g(z) + \delta^{-\frac{1}{2}} \|g\|_{L^2(\mathbb{C})} \right).$$

After choosing $\delta = V^{-1}(z)$ in the above, we further get

$$I_{\frac{1}{2}} g(z) \leq C \left(V(z)^{-\frac{1}{2}} M g(z) + V(z)^{\frac{1}{2}} \|g\|_{L^2(\mathbb{C})} \right).$$

Square both sides of the above inequality, multiply them by V , and then integrate over \mathbb{C} . One has

$$(2.2) \quad \left(\int_{\mathbb{C}} \left| I_{\frac{1}{2}} g(z) \right|^2 V(z) dv_z \right)^{\frac{1}{2}} \leq C (\|Mg\|_{L^2(\mathbb{C})} + \|V\|_{L^2(\mathbb{C})} \|g\|_{L^2(\mathbb{C})}) \\ \leq C (1 + \|V\|_{L^2(\mathbb{C})}) \|g\|_{L^2(\mathbb{C})}.$$

In the last inequality, we have used the boundedness of the Maximal function operator in $L^2(\mathbb{C})$ space.

We further employ the standard rescaling technique to get rid of the constant 1 in the last line of (2.2). Indeed, for any $k > 0$, replacing V by kV in (2.2) and then dividing by $k^{\frac{1}{2}}$, we have

$$\left(\int_{\mathbb{C}} \left| I_{\frac{1}{2}} g(z) \right|^2 V(z) dv_z \right)^{\frac{1}{2}} \leq C \left(\frac{1}{k^{\frac{1}{2}}} + k^{\frac{1}{2}} \|V\|_{L^2(\mathbb{C})} \right) \|g\|_{L^2(\mathbb{C})}.$$

Choosing $k = \frac{1}{\|V\|_{L^2(\mathbb{C})}}$, this then completes the proof of (2.1).

Therefore if V is $L^2(\mathbb{C})$, $I_{\frac{1}{2}}$ becomes a bounded operator from $L^2(\mathbb{C})$ to $L^2_V(\mathbb{C})$. Consequently, let $I_{\frac{1}{2}}^*$ be the dual of $I_{\frac{1}{2}}$ with respect to the following inner product

$$\langle f, g \rangle := \int_{\mathbb{C}} f(z) \overline{g(z)} dv_z \quad \text{for all } f, g \in L^2(\mathbb{C}).$$

Then $I_{\frac{1}{2}}^*$ is bounded from $L^2_{V^{-1}}(\mathbb{C})$ to $L^2(\mathbb{C})$ satisfying for any $f \in L^2_{V^{-1}}(\mathbb{C})$,

$$\|I_{\frac{1}{2}}^* f(z)\|_{L^2(\mathbb{C})} \leq C \|V\|_{L^2(\mathbb{C})}^{\frac{1}{2}} \|f\|_{L^2_{V^{-1}}(\mathbb{C})}.$$

Note that I_{α} is a convolution operator. So one has $I_{\frac{1}{2}}^* = I_{\frac{1}{2}}$. Making use of the semi-group property of the fractional integrals, we obtain $I_1 = I_{\frac{1}{2}} \circ I_{\frac{1}{2}}^*$ is a bounded operator from $L^2_{V^{-1}}(\mathbb{C})$ into $L^2_V(\mathbb{C})$ with the desired estimate. \square

3. Proof of Theorem 1.1

Throughout this section, we restrict on the case when $n = 1$. The key ingredient of Chanillo and Sawyer's proof in [3] lies in a pointwise inequality in the spirit of a technical result of Sawyer in [11], together with a weighted inequality for the Riesz integral operator I_1 . The proof of our Theorem 1.1 essentially follows their idea. Instead of using a Sawyer inequality, we only need the following trivial identity (3.1) for the Cauchy kernel.

$$(3.1) \quad \frac{1}{z - \zeta} + \sum_{l=0}^{m-1} \frac{z^l}{\zeta^{l+1}} = \frac{z^m}{\zeta^m(z - \zeta)}, \quad \text{for all } \zeta \neq z \text{ nor } 0.$$

As usual, denote by D_R the disk in \mathbb{C} centered at 0 with radius R .

Lemma 3.1. *Let $u \in H^1(\mathbb{C})$ has compact support. Then for almost every $z \in \mathbb{C}$,*

$$(3.2) \quad u(z) = \frac{1}{\pi} \int_{\mathbb{C}} \frac{\bar{\partial}u(\zeta)}{z - \zeta} dv_{\zeta}.$$

If in addition that u vanishes near 0, then for any $l \geq 0$,

$$(3.3) \quad \int_{\mathbb{C}} \frac{\bar{\partial}u(\zeta)}{\zeta^{l+1}} dv_{\zeta} = 0.$$

Proof. Let $R_0 > 0$ be such that $\text{supp } u \subset D_{R_0}$. For each $z \in \mathbb{C}$, pick $R > R_0$ so that $z \in D_R$. If $u \in C^1(\mathbb{C})$, by the Cauchy-Green formula (see for instance [13]),

$$u(z) = \frac{1}{2\pi i} \int_{\partial D_R} \frac{u(\zeta)}{\zeta - z} d\zeta - \frac{1}{\pi} \int_{D_R} \frac{\bar{\partial}u(\zeta)}{\zeta - z} dv_{\zeta} = \frac{1}{\pi} \int_{\mathbb{C}} \frac{\bar{\partial}u(\zeta)}{z - \zeta} dv_{\zeta}.$$

Here we used the fact that $\text{supp } u \subset D_{R_0}$. For general u in $H_0^1(D_{R_0})$, letting $u_j \in C_c^1(D_{R_0}) \rightarrow u$ in $H^1(D_{R_0})$ norm, then for any $R > R_0$,

$$\begin{aligned} & \left\| u - \frac{1}{\pi} \int_{\mathbb{C}} \frac{\bar{\partial}u(\zeta)}{\cdot - \zeta} dv_{\zeta} \right\|_{L^2(D_R)} \\ & \leq \|u - u_j\|_{L^2(D_R)} + \frac{1}{\pi} \left\| \int_{D_R} \frac{\bar{\partial}(u(\zeta) - u_j(\zeta))}{\cdot - \zeta} dv_{\zeta} \right\|_{L^2(D_R)} \\ & \leq C \|u - u_j\|_{H^1(D_R)} \\ & = C \|u - u_j\|_{H^1(D_{R_0})} \rightarrow 0 \end{aligned}$$

as $j \rightarrow \infty$. Here the validity of the second inequality was based on the boundedness of the solid Cauchy integral operator

$$Tf := -\frac{1}{\pi} \int_{D_R} \frac{f(\zeta)}{z - \zeta} dv_{\zeta}$$

from $L^2(D_R)$ space to itself. This proves (3.2).

When u vanishes near 0, the integral (3.3) is well defined. Moreover, $\int_{\mathbb{C}} \frac{\bar{\partial}u(\zeta)}{z - \zeta} dv_{\zeta}$ is zero near 0 by (3.2). Thus

$$\int_{\mathbb{C}} \frac{\bar{\partial}u(\zeta)}{\zeta^{l+1}} dv_{\zeta} = -\frac{1}{l!} \frac{\partial^l}{\partial z^l} \int_{\mathbb{C}} \frac{\bar{\partial}u(\zeta)}{z - \zeta} dv_{\zeta} \Big|_{z=0} = 0.$$

□

Proof of Theorem 1.1: Without loss of generality, we assume $z_0 = 0$. Let χ_B be the characteristic function for a set B . Fix an $r > 0$ small enough such that $D_{2r} \subset \Omega$, $\|u\|_{H^1(D_{2r})}$ is finite, and

$$\left\| V\chi_{D_r} + \frac{r}{1 + |z|^2} \right\|_{L^2(\mathbb{C})}^2 \leq \frac{\pi^2}{2C_0},$$

where C_0 is the universal constant in Theorem 2.2. Replacing V by $V\chi_{D_r} + \frac{r}{1+|z|^2}$, still denoted by V , we have (1.2) holds on D_{2r} with $V \in L^2(\mathbb{C})$,

$$(3.4) \quad V > 0 \text{ on } \mathbb{C}; \quad V \geq C_r \text{ on } D_{2r}$$

for some positive constant C_r dependent only on r , and

$$(3.5) \quad \|V\|_{L^2(\mathbb{C})}^2 \leq \frac{\pi^2}{2C_0}.$$

We shall show that $u = 0$ on $D_{\frac{r}{2}}$. Then the rest of the proof for the strong unique continuation property follows from a standard propagation argument.

Choose $\eta \in C_c^\infty(\mathbb{C})$ such that $\eta = 1$ on D_r ; $0 \leq \eta \leq 1$ and $|\nabla\eta| \leq \frac{2}{r}$ on $D_{2r} \setminus D_r$; $\eta = 0$ outside D_{2r} . Let $\psi \in C^\infty(\mathbb{C})$ be such that $\psi = 0$ in D_1 ; $0 \leq \psi \leq 1$ and $|\nabla\psi| \leq 2$ on $D_2 \setminus D_1$; $\psi = 1$ outside D_2 . For each $k \geq \frac{4}{r}$ (then $\frac{2}{k} \leq \frac{r}{2}$), let $\psi_k(z) = \psi(kz)$, $z \in \mathbb{C}$. Note that $\psi_k\eta u \in H^1(\mathbb{C})$ and is supported inside $D_{2r} \setminus D_{\frac{1}{k}}$. We apply Lemma 3.1 to $\psi_k\eta u$ and obtain for $z \in D_r$, $m \in \mathbb{Z}^+$,

$$\begin{aligned} |\psi_k(z)|^2 |u(z)|^2 &= \frac{1}{\pi^2} \left| \int_{\mathbb{C}} \frac{\bar{\partial}(\psi_k(\zeta)\eta(\zeta)u(\zeta))}{z - \zeta} dv_\zeta \right|^2 \\ &= \frac{1}{\pi^2} \left| \int_{\mathbb{C}} \left(\frac{1}{z - \zeta} + \sum_{l=0}^{m-1} \frac{z^l}{\zeta^{l+1}} \right) \bar{\partial}(\psi_k(\zeta)\eta(\zeta)u(\zeta)) dv_\zeta \right|^2. \end{aligned}$$

Making use of (3.1), we have

$$\begin{aligned} &\int_{D_r} \frac{|\psi_k(z)|^2 |u(z)|^2}{|z|^{2m}} V(z) dv_z \\ &= \frac{1}{\pi^2} \int_{D_r} \frac{1}{|z|^{2m}} \left| \int_{\mathbb{C}} \left(\frac{1}{z - \zeta} + \sum_{l=0}^{m-1} \frac{z^l}{\zeta^{l+1}} \right) \bar{\partial}(\psi_k(\zeta)\eta(\zeta)u(\zeta)) dv_\zeta \right|^2 V(z) dv_z \\ &\leq \frac{1}{\pi^2} \int_{D_r} \left(\int_{\mathbb{C}} \frac{1}{|z - \zeta|} \frac{|\bar{\partial}(\psi_k(\zeta)\eta(\zeta)u(\zeta))|}{|\zeta|^m} dv_\zeta \right)^2 V(z) dv_z \\ &\leq \frac{1}{\pi^2} \left\| I_1 \left(\frac{|\bar{\partial}(\psi_k\eta u)|}{|\cdot|^m} \right) \right\|_{L_V^2(\mathbb{C})}^2. \end{aligned}$$

Applying Theorem 2.2 and (3.5), we further obtain

$$\begin{aligned}
& \int_{D_r} \frac{|\psi_k(z)|^2 |u(z)|^2}{|z|^{2m}} V(z) dv_z \\
& \leq \frac{C_0}{\pi^2} \|V\|_{L^2(\mathbb{C})}^2 \int_{\mathbb{C}} \frac{|\bar{\partial}(\psi_k(z)\eta(z)u(z))|^2}{|z|^{2m}} \frac{dv_z}{V(z)} \\
(3.6) \quad & \leq \frac{1}{2} \left(\int_{\mathbb{C}} \frac{|\bar{\partial}\psi_k(z)|^2 |u(z)|^2}{|z|^{2m} V(z)} dv_z + \int_{D_r} \frac{|\psi_k(z)|^2 |\bar{\partial}u(z)|^2}{|z|^{2m} V(z)} dv_z + \right. \\
& \quad \left. + \int_{D_{2r} \setminus D_r} \frac{|\bar{\partial}(\eta(z)u(z))|^2}{|z|^{2m} V(z)} dv_z \right) \\
& =: \frac{A}{2} + \frac{B}{2} + \frac{C}{2}.
\end{aligned}$$

We first have $\lim_{k \rightarrow \infty} A = 0$. Indeed, since $\bar{\partial}\psi_k$ is only supported on $D_{\frac{2}{k}} \setminus D_{\frac{1}{k}}$, and $V > C_r$ on $D_{\frac{2}{k}} (\subset D_r)$,

$$A \leq \int_{\frac{1}{k} < |z| < \frac{2}{k}} \frac{|\nabla\psi_k(z)|^2 |u(z)|^2}{|z|^{2m} V(z)} dv_z \leq \frac{k^{2m+2}}{C_r} \int_{|z| < \frac{2}{k}} |u(z)|^2 dv_z \rightarrow 0$$

as $k \rightarrow \infty$, as a consequence of the fact that u vanishes to infinite order at 0.

On the other hand, applying the inequality (1.2) to B , we get

$$\frac{B}{2} \leq \frac{1}{2} \int_{D_r} \frac{|\psi_k(z)|^2 |u(z)|^2}{|z|^{2m}} V(z) dv_z.$$

Subtract $\frac{B}{2}$ from both sides of (3.6) and let $k \rightarrow \infty$. Then for each $m \in \mathbb{Z}^+$, by Fatou's Lemma,

$$\begin{aligned}
\int_{D_r} \frac{|u(z)|^2}{|z|^{2m}} V(z) dv_z & \leq \lim_{k \rightarrow \infty} \int_{D_r} \frac{|\psi_k(z)|^2 |u(z)|^2}{|z|^{2m}} V(z) dv_z \\
& \leq \int_{D_{2r} \setminus D_r} \frac{|\bar{\partial}(\eta(z)u(z))|^2}{|z|^{2m} V(z)} dv_z.
\end{aligned}$$

Now multiplying both sides of the above inequalities by r^{2m} , then

$$\begin{aligned}
(3.7) \quad \int_{D_r} \frac{r^{2m}}{|z|^{2m}} |u(z)|^2 V(z) dv_z & \leq \int_{D_{2r} \setminus D_r} \frac{r^{2m} |\bar{\partial}(\eta(z)u(z))|^2}{|z|^{2m} V(z)} dv_z \\
& \leq \int_{D_{2r} \setminus D_r} \frac{|\bar{\partial}(\eta(z)u(z))|^2}{V(z)} dv_z.
\end{aligned}$$

Shrinking the integral domain of the left hand side of (3.7) to $D_{\frac{r}{2}}$, and making use of (3.4), we further infer

$$2^{2m} \int_{D_{\frac{r}{2}}} |u(z)|^2 V(z) dv_z \leq \frac{1}{C_r} \int_{D_{2r}} |\bar{\partial}(\eta(z)u(z))|^2 dv_z = \tilde{C}_r \|u\|_{H^1(D_{2r})}^2,$$

for some constant \tilde{C}_r dependent only on r . Lastly, letting $m \rightarrow \infty$, we have $u = 0$ on $D_{\frac{r}{2}}$. \square

It is worth pointing out that when $V \in L^p, p > 2$, the weighted Hardy-Littlewood-Sobolev inequality is not needed to obtain the unique continuation property for (1.2). Indeed, the strong unique continuation property follows by repeating the same proof as in Theorem 1.1, except with those integrals over the domain \mathbb{C} substituted by those over $D_{2r}, r < \frac{1}{2}$, and with Theorem 2.2 replaced by the following inequality

$$(3.8) \quad \|I_1 f\|_{L^2_V(D_1)} \leq C_p \|V\|_{L^p(D_1)} \|f\|_{L^2_{V^{-1}}(D_1)} \quad \text{for all } f \in L^2_{V^{-1}}(D_1)$$

for a constant C_p dependent only on p . (3.8) can be proved by simply using Hölder inequality. This is because when $p > 2$, $\|\frac{1}{\cdot - z}\|_{L^q(D_1)}$ is uniformly bounded on D_1 by some constant C_p , where q is the dual of p , i.e., $\frac{1}{p} + \frac{1}{q} = 1$. Therefore,

$$\begin{aligned} \|I_1 f\|_{L^2_V(D_1)}^2 &\leq \int_{D_1} \left| \int_{D_1} \frac{|f(\zeta)|}{|\zeta - z|^{\frac{1}{2}} V(\zeta)^{\frac{1}{2}}} \cdot \frac{V(\zeta)^{\frac{1}{2}}}{|\zeta - z|^{\frac{1}{2}}} dv_\zeta \right|^2 V(z) dv_z \\ &\leq \int_{D_1} \int_{D_1} \frac{|f(\zeta)|^2}{|\zeta - z| V(\zeta)} dv_\zeta \cdot \int_{D_1} \frac{V(\zeta)}{|\zeta - z|} dv_\zeta V(z) dv_z \\ &\leq C_p \|V\|_{L^p(D_1)} \int_{D_1} \int_{D_1} \frac{|f(\zeta)|^2}{|\zeta - z| V(\zeta)} dv_\zeta V(z) dv_z \\ &= C_p \|V\|_{L^p(D_1)} \int_{D_1} \frac{|f(\zeta)|^2}{V(\zeta)} \int_{D_1} \frac{V(z)}{|\zeta - z|} dv_z dv_\zeta \\ &\leq C_p^2 \|V\|_{L^p(D_1)}^2 \|f\|_{L^2_{V^{-1}}(D_1)}^2. \end{aligned}$$

Clearly the above approach no longer works when, for instance, the potential V is exactly L^2 (namely, in L^2 but not in L^p for any $p > 2$) as in the following example. For those cases, we have to use Theorem 1.1 to justify the nonexistence of nontrivial flat solutions.

Example 1. For $0 < \epsilon < \frac{1}{2}$, let

$$u_0(z) = e^{-(-\ln|z|)^\epsilon} \quad \text{and} \quad V(z) = \frac{\epsilon}{2|z|(-\ln|z|)^{1-\epsilon}}$$

on $D_{\frac{1}{2}}$. Then $V \in L^2(D_{\frac{1}{2}})$, yet $V \notin L^p(D_{\frac{1}{2}})$ for any $p > 2$. By Theorem 1.1, the equation $|\bar{\partial}u| = V|u|$ has no nontrivial flat solution. Since u_0 is continuous on $D_{\frac{1}{2}}$ and satisfies $|\bar{\partial}u| = V|u|$ in $D_{\frac{1}{2}} \setminus \{0\}$ pointwisely, by Lemma A.1 we have $u_0 \in H^1(D_{\frac{1}{2}})$ and satisfies $|\bar{\partial}u| = V|u|$ on $D_{\frac{1}{2}}$. Thus u_0 can not be flat in particular. On the other hand, one can directly verify that $u_0^{-1}(0) = \{0\}$, yet u_0 fails to vanish to infinite order (in fact, (1.1) fails for all $m \geq 3$) at 0.

When $V \in L^p, p < 2$, the strong unique continuation property fails in general as seen below.

Example 2. For each $0 < p < 2$, choose $\epsilon \in (0, \frac{2-p}{p})$ (so that $(\epsilon + 1)p < 2$). Then $u_\epsilon(z) := e^{-\frac{1}{|z|^\epsilon}}$ vanishes to infinite order at 0 and satisfies $|\bar{\partial}u| = V|u|$ with $V = \frac{\epsilon}{2|z|^{\epsilon+1}} \in L^p(D_1)$ on \mathbb{C} .

More strikingly, Mandache constructed an example in [8] with an $L^p, p < 2$ potential, where even the weak unique continuation property fails.

Example 3. [8] There exist two functions $u \in C_c^\infty(\mathbb{C})$ and $V \in L^p(\mathbb{C}), 0 < p < 2$, such that u is supported on D_1 and satisfies $\bar{\partial}u = Vu$.

Despite the above examples in $p < 2$, the first author showed (in Lemma 7 [9]) that the strong unique continuation property may still be expected if the target dimension $N = 1$ and V takes certain special form, such as a multiple of $\frac{1}{|z|}$. Note that $\frac{1}{|z|} \notin L^2$. It should be noted that the strong unique continuation property no longer holds for this potential when $N = 2$. In fact, Wolff and the first author in [10] constructed a smooth function $v_0 : \mathbb{C} \rightarrow \mathbb{C}$ which vanishes to infinite order at 0 and satisfies $|\Delta v| \leq \frac{C}{|z|}|\nabla v|$ for some constant $C > 0$. (See also [1] by Alinhac and Baouendi for another example in the same setting.) Letting $u_0 := (\partial\bar{\mathcal{R}}v_0, \partial\mathfrak{I}v_0)$, then $u_0 : \mathbb{C} \rightarrow \mathbb{C}^2$ vanishes to infinite order at 0 and satisfies $|\bar{\partial}u| \leq \frac{C}{|z|}|u|$ for some constant C .

4. Proof of Theorem 1.3 and Corollary 1.4

We shall assume $n \geq 2$ in this section. Note that the inequality (1.2) reads as

$$(4.1) \quad |\bar{\partial}u| := \left(\sum_{j=1}^n \sum_{k=1}^N |\bar{\partial}_j u_k|^2 \right)^{\frac{1}{2}} \leq V \left(\sum_{k=1}^N |u_k|^2 \right)^{\frac{1}{2}} := V|u|.$$

Proof of Theorem 1.3: Without loss of generality, assume $n = 2$. We only need to show that for any $r_2 \geq r_1 > 0, s > 0$, if u satisfies (1.2) on the bidisk $D_{r_2} \times D_s$ and $u = 0$ on $D_{r_1} \times D_s$, then $u = 0$ on $D_{r_2} \times D_s$.

Since $V \in L_{loc}^2(D_{r_2} \times D_s)$, by Fubini's theorem, for almost every $z_2 \in D_s$, $V(\cdot, z_2) \in L_{loc}^2(D_{r_2})$, and similarly for $u(\cdot, z_2) \in H_{loc}^1(D_{r_2})$. Restricting (4.1) at each such $z_2 = c_2$, we have $v := u(\cdot, c_2)$ on D_{r_2} satisfies

$$|\bar{\partial}v| = \left(\sum_{k=1}^N |\bar{\partial}_1 u_k(\cdot, c_2)|^2 \right)^{\frac{1}{2}} \leq |\bar{\partial}u(\cdot, c_2)| \leq V(\cdot, c_2)|u(\cdot, c_2)| = V(\cdot, c_2)|v|.$$

Since $v = 0$ on D_{r_1} , applying Theorem 1.1 we have $v = 0$ on D_{r_2} . Thus $u = 0$ on $D_{r_2} \times D_s$. The proof is complete. \square

Proof of Corollary 1.4: Let $u := f - g$. Then $u \in H_{loc}^1(\Omega)$ vanishes in an open subset and satisfies

$$\bar{\partial}u = Au \quad \text{on } \Omega.$$

Let $V := \sqrt{\sum_{j,k=1}^N |A_{jk}|^2}$, the matrix norm of A . Then u satisfies $|\bar{\partial}u| \leq V|u|$ with $V \in L_{loc}^2(\Omega)$. By Theorem 1.3, we have $u \equiv 0$ on Ω . \square

One may modify the one-dimensional Example 1 to obtain an inequality in higher dimensions with an exact L^2 potential, where Theorem 1.3 can be applied to get the weak continuation property.

Example 4. For $0 < \epsilon < \frac{1}{2}$, let

$$u_0(z_1, z_2) = e^{-(\ln|z_1|)^\epsilon} e^{-(\ln|z_2|)^\epsilon}$$

and

$$V(z_1, z_2) = \frac{\epsilon}{2} \left(\frac{1}{|z_1|^2 (-\ln|z_1|)^{2-2\epsilon}} + \frac{1}{|z_2|^2 (-\ln|z_2|)^{2-2\epsilon}} \right)^{\frac{1}{2}}$$

on $D_{\frac{1}{2}} \times D_{\frac{1}{2}}$. Then $V \in L^2(D_{\frac{1}{2}} \times D_{\frac{1}{2}})$, yet $V \notin L^p(D_{\frac{1}{2}} \times D_{\frac{1}{2}})$ for any $p > 2$. Moreover, since u_0 is continuous on $D_{\frac{1}{2}} \times D_{\frac{1}{2}}$ and satisfies $|\bar{\partial}u| = V|u|$ in $(D_{\frac{1}{2}} \setminus \{0\}) \times (D_{\frac{1}{2}} \setminus \{0\})$, $u_0 \in H^1(D_{\frac{1}{2}} \times D_{\frac{1}{2}})$ and satisfies $|\bar{\partial}u| = V|u|$ on $D_{\frac{1}{2}} \times D_{\frac{1}{2}}$ by Corollary A.2. By Theorem 1.3, any nontrivial solution, and thus u_0 , can not vanish in open subsets of $D_{\frac{1}{2}} \times D_{\frac{1}{2}}$. In fact, one can directly verify that u_0 only vanishes on $\{z_1 = 0\} \cup \{z_2 = 0\}$ in $D_{\frac{1}{2}} \times D_{\frac{1}{2}}$.

Similarly, making use of Mandache's example, one can construct an example in $n \geq 2$ where the weak unique continuation property fails if the potential is at most $L^p, p < 2$.

Example 5. For each $0 < p < 2$, there exist nontrivial $u \in C_c^\infty(\mathbb{C}^2)$, $V \in L^p(\mathbb{C}^2)$ such that u is supported on $D_1 \times D_1$ and satisfies $|\bar{\partial}u| \leq V|u|$ on \mathbb{C}^2 .

Proof. Let w, W be as in Mandache's Example 3 with $w \in C_c^\infty(D_1)$, $W \in L^p(\mathbb{C})$ and $|\bar{\partial}w| \leq W|w|$ on \mathbb{C} . Define $u(z_1, z_2) := w(z_1)w(z_2)$, $(z_1, z_2) \in \mathbb{C}^2$. Then $u \in C_c^\infty(\mathbb{C}^2)$ with support in $D_1 \times D_1$. One can check that for $(z_1, z_2) \in \mathbb{C}^2$,

$$\begin{aligned} |\bar{\partial}u(z_1, z_2)| &\leq |\bar{\partial}_1 w(z_1)| |w(z_2)| + |w(z_1)| |\bar{\partial}_2 w(z_2)| \\ &\leq W(z_1) |w(z_1)| |w(z_2)| + W(z_2) |w(z_1)| |w(z_2)| \\ &= (W(z_1)\chi_{D_1}(z_2) + W(z_2)\chi_{D_1}(z_1)) |u(z_1, z_2)|. \end{aligned}$$

Letting $V(z_1, z_2) := W(z_1)\chi_{D_1}(z_2) + W(z_2)\chi_{D_1}(z_1)$. Then $|\bar{\partial}u| \leq V|u|$ with

$$\|V\|_{L^p(\mathbb{C}^2)}^p \leq C \left(\int_{\mathbb{C}} |W(z_1)|^p dv_{z_1} + \int_{\mathbb{C}} |W(z_2)|^p dv_{z_2} \right) < \infty$$

for a constant $C > 0$. \square

The following theorem gives an example where the strong unique continuation property may hold when the potential V is a multiple of $\frac{1}{|z|}$ as below. We note that when $n \geq 2$, $\frac{1}{|z|}$ is in L^2 but not in L^{2n} . Let B_1 be the unit ball centered at $0 \in \mathbb{C}^n$.

Theorem 4.1. *Let u be a smooth function from $B_1 \subset \mathbb{C}^n$ to \mathbb{C} satisfying*

$$(4.2) \quad |\bar{\partial}u| \leq \frac{C}{|z|}|u|$$

for some constant $C > 0$. If u vanishes to infinite order at 0 , then u vanishes identically on B_1 .

Proof. For each $z \in B_1 \setminus \{0\}$, define $w(\tau) := u(\tau z)$, $\tau \in D_{\frac{1}{|z|}}$. Then w vanishes to infinite order at 0 and satisfies

$$|\bar{\partial}w(\tau)| \leq |z| |\bar{\partial}u(\tau z)| \leq \frac{C}{|\tau|} |w(\tau)|.$$

Making use of Lemma 7 in [9], we have w vanishes on $D_{\frac{1}{|z|}}$, and thus u vanishes identically on B_1 . \square

Appendix A.

The following lemma is known in folklore, and is particularly useful while checking for weak derivatives in examples. For convenience of the reader, we provide a proof below.

Lemma A.1. *Let $f : D_1 \rightarrow \mathbb{C}$ and $f \in L^1(D_1)$. Suppose $u : D_1 \rightarrow \mathbb{C}$ with $u \in L^2(D_1)$ satisfies*

$$\bar{\partial}u = f \text{ in } D_1 \setminus \{0\}$$

in weak sense. Then

$$\bar{\partial}u = f \text{ in } D_1$$

in weak sense.

Proof. Let $\eta \in C_c^\infty(D_1)$ be a cut-off function with $\eta = 1$ on $D_{\frac{1}{2}}$ and $|\nabla\eta| < 3$ on D_1 . For each $k > 0$, let $\eta_k(z) = \eta(kz)$, $z \in \mathbb{C}$. Given a fixed testing function $\phi \in C_c^\infty(D_1)$,

$$\begin{aligned} & \int_{D_1} u(z) \bar{\partial}\phi(z) dv_z \\ &= \int_{D_1} u(z) \bar{\partial}(\eta_k(z)\phi(z)) dv_z + \int_{D_1} u(z) \bar{\partial}((1 - \eta_k(z))\phi(z)) dv_z := A + B. \end{aligned}$$

Since $(1 - \eta_k)\phi$ is a testing function on $D_1 \setminus \{0\}$, by assumption

$$B = - \int_{D_1} f(z) (1 - \eta_k(z)) \phi(z) dv_z \rightarrow - \int_{D_1} f(z) \phi(z) dv_z$$

when $k \rightarrow \infty$. For A , since η_k and $|\bar{\partial}\eta_k|$ are only supported on $D_{\frac{1}{k}}$ with $|\eta_k| \leq 1$ and $|\bar{\partial}\eta_k| \leq 3k$, we have by Hölder inequality

$$\begin{aligned} |A| &\leq \int_{D_1} |u(z)| |\bar{\partial}\eta_k(z)| |\phi(z)| dv_z + \int_{D_1} |u(z)| |\eta_k(z)| |\bar{\partial}\phi(z)| dv_z \\ &\leq 3k \int_{D_{\frac{1}{k}}} |u(z)\phi(z)| dv_z + \int_{D_{\frac{1}{k}}} |u(z)\bar{\partial}\phi(z)| dv_z \\ &\leq 3k \frac{\sqrt{\pi}}{k} \|u\phi\|_{L^2(D_{\frac{1}{k}})} + \frac{\sqrt{\pi}}{k} \|u\bar{\partial}\phi\|_{L^2(D_{\frac{1}{k}})} \\ &\leq C \|u\|_{L^2(D_{\frac{1}{k}})} \end{aligned}$$

for some constant $C > 0$ (dependent only on ϕ). Letting $k \rightarrow \infty$, we get $|A| \rightarrow 0$. The proof is complete. \square

We remark that the assumption $u \in L^2(D_1)$ in Lemma A.1 can not be further relaxed, as indicated by the following example.

Example 6. Let $u_0(z) = \frac{1}{z}$, $z \in D_1$. Then u_0 satisfies

$$\bar{\partial}u = 0 \quad \text{in } D_1 \setminus \{0\}.$$

However, u_0 is not a weak solution to $\bar{\partial}u = 0$ in D_1 .

Proof. We only need to verify for a general $\phi \in C_c^\infty(D_1)$, $\int_{D_1} \frac{\bar{\partial}\phi(z)}{z} dv_z \neq 0$. Indeed, by Stokes' theorem,

$$\begin{aligned} \int_{D_1} \frac{\bar{\partial}\phi(z)}{z} dv_z &= \lim_{\epsilon \rightarrow 0} \int_{D_1 \setminus D_\epsilon} \frac{\bar{\partial}\phi(z)}{z} dv_z \\ &= \frac{1}{2} \lim_{\epsilon \rightarrow 0} \int_{D_1 \setminus D_\epsilon} \bar{\partial} \left(\frac{\phi(z)}{z} dz \right) \\ &= - \frac{1}{2} \lim_{\epsilon \rightarrow 0} \int_{\partial D_\epsilon} \frac{\phi(z)}{z} dz \\ &= - \phi(0) \pi i. \end{aligned}$$

\square

One can immediately extend Lemma A.1 to higher dimensions as follows.

Corollary A.2. *Let $f_j : D_1 \rightarrow \mathbb{C}$ and $f_j \in L^1(D_1)$, $j = 1, 2$. Suppose $u_j : D_1 \rightarrow \mathbb{C}$ with $u_j \in L^2(D_1)$ and satisfies*

$$\bar{\partial}u_j = f_j \text{ in } D_1 \setminus \{0\}$$

in weak sense for each j . Then $u(z_1, z_2) := u_1(z_1)u_2(z_2)$ satisfies

$$\bar{\partial}_1 u(z_1, z_2) = f_1(z_1)u_2(z_2) \text{ and } \bar{\partial}_2 u(z_1, z_2) = u_1(z_1)f_2(z_2) \text{ in } D_1 \times D_1$$

in weak sense.

Proof. We only prove the first equation holds in $D_1 \times D_1$ in weak sense. For a fixed testing function $\phi \in C_c^\infty(D_1 \times D_1)$, by Fubini's theorem,

$$\int_{D_1 \times D_1} u(z_1, z_2) \bar{\partial}_1(\phi(z_1, z_2)) dv_z = \int_{D_1} u_2(z_2) \int_{D_1} u_1(z_1) \bar{\partial}_1(\phi(z_1, z_2)) dv_{z_1} dv_{z_2}.$$

Since $\phi(\cdot, z_2) \in C_c^\infty(D_1)$ for each $z_2 \in D_1$, applying Lemma A.1 to u_1 , we get

$$\begin{aligned} \int_{D_1 \times D_1} u(z_1, z_2) \bar{\partial}_1(\phi(z_1, z_2)) dv_z &= - \int_{D_1} u_2(z_2) \int_{D_1} f_1(z_1) \phi(z_1, z_2) dv_{z_1} dv_{z_2} \\ &= - \int_{D_1 \times D_1} f_1(z_1) u_2(z_2) \phi(z_1, z_2) dv_z. \end{aligned}$$

□

References

- [1] ALINHAC, S.; BAOUENDI, S. A counterexample to strong uniqueness for partial differential equations of Schrödinger's type. *Comm. Partial Differential Equations* **19** (1994), no. 9-10, 1727–1733. [MR1294476](#), [Zbl 0806.35023](#), doi: [10.1080/03605309408821069](#).
- [2] BELL, S.; LEMPert, L. A C^∞ Schwarz reflection principle in one and several complex variables. *J. Differential Geom.* **32** (1990), no. 3, 899–915. [MR1078165](#), [Zbl 0716.32002](#), doi: [10.4310/jdg/1214445538](#).
- [3] CHANILLO, S.; SAWYER, E. Unique continuation for $\Delta + v$ and the C. Fefferman-Phong class. *Trans. Amer. Math. Soc.* **318** (1990), no. 1, 275–300. [MR0958886](#), [Zbl 0702.35034](#), doi: [10.2307/2001239](#).
- [4] CHANILLO, S.; WHEEDEN, R. L^p -estimates for fractional integrals and Sobolev inequalities with applications to Schrödinger operators. *Comm. Partial Differential Equations* **10** (1985), no. 9, 1077–1116. [MR0806256](#), [Zbl 0578.46024](#), doi: [10.1080/03605308508820401](#).
- [5] FASSINA, M.; PAN, Y. A local obstruction for elliptic operators with real analytic coefficients on flat germs. *Ann. Fac. Sci. Toulouse Math.* (6) **31** (2022), no. 5, 1343–1363. [MR4520802](#), [Zbl 7632688](#), doi: [10.5802/afst.1722](#).
- [6] KENIG, C.; WANG, J.-N.. Quantitative uniqueness estimates for second order elliptic equations with unbounded drift. *Math. Res. Lett.* **22** (2015), no. 4, 1159–1175. [MR3391881](#), [Zbl 1326.35098](#), doi: [10.4310/MRL.2015.v22.n4.a10](#).
- [7] LIU, Y.; PAN, Y.; ZHANG, Y. A derivative formula for the solid Cauchy integral operator and its applications. *Integral Equations Operator Theory* **94** (2022), no. 4, Paper No. 38, 23 pp. [MR4502708](#), [Zbl 7615125](#), doi: [10.1007/s00020-022-02717-0](#).

- [8] MANDACHE, N. A counterexample to unique continuation in dimension two. *Comm. Anal. Geom.* **10** (2002), no. 1, 1–10. [MR1894138](#), [Zbl 1001.35019](#), doi: [10.4310/CAG.2002.v10.n1.a1](#).
- [9] PAN, Y. Unique continuation for Schrödinger operators with singular potentials. *Comm. Partial Differential Equations* **17** (1992), no. 5-6, 953–965. [MR1177300](#), [Zbl 0810.35005](#), doi: [10.1080/03605309208820871](#).
- [10] PAN, Y.; WOLFF, T. A remark on unique continuation. *J. Geom. Anal.* **8** (1998), no. 4, 599–604. [MR1724207](#), [Zbl 0960.35012](#), doi: [10.1007/BF02921714](#).
- [11] SAWYER, E. Unique continuation for Schrödinger operators in dimension three or less. *Ann. Inst. Fourier (Grenoble)* **34** (1984), no. 3, 189–200. [MR0762698](#), [Zbl 0535.35007](#), doi: [10.5802/aif.982](#).
- [12] SEO, I. A remark on unique continuation for the Cauchy-Riemann operator. *Bull. Korean Math. Soc.* **52** (2015), no. 5, 1753–1757. [MR3406034](#), [Zbl 1329.35088](#), doi: [10.4134/BKMS.2015.52.5.1753](#).
- [13] VEKUA, I. N. *Generalized analytic functions. International Series of Monographs on Pure and Applied Mathematics* Vol. 25, Pergamon Press, 1962 xxix+668 pp. [MR0150320](#), [Zbl 0100.07603](#).
- [14] WOLFF, T. A counterexample in a unique continuation problem. *Comm. Anal. Geom.* **2** (1994), no. 1, 79–102. [MR1312679](#), [Zbl 0836.35023](#), doi: [10.4310/CAG.1994.v2.n1.a5](#).

DEPARTMENT OF MATHEMATICAL SCIENCES, PURDUE UNIVERSITY FORT WAYNE, FORT WAYNE, IN 46805-1499, USA.
pan@pfw.edu

DEPARTMENT OF MATHEMATICAL SCIENCES, PURDUE UNIVERSITY FORT WAYNE, FORT WAYNE, IN 46805-1499, USA.
zhangyu@pfw.edu